

CHEREDNIK ALGEBRAS AND DIFFERENTIAL OPERATORS ON QUASI-INVARIANTS

YURI BEREST, PAVEL ETINGOF AND VICTOR GINZBURG

Abstract

We develop representation theory of the rational Cherednik algebra H_c associated to a finite Coxeter group W in a vector space $\mathfrak{h}$, and a parameter ‘ c ’. We use it to show that, for integral values of ‘ c ’, the algebra H_c is simple and Morita equivalent to $\mathcal{D}(\mathfrak{h})\#W$, the cross product of W with the algebra of polynomial differential operators on $\mathfrak{h}$.

Chalykh, Feigin, and Veselov [CV], [FV], introduced an algebra, Q_c , of *quasi-invariant* polynomials on $\mathfrak{h}$, such that $\mathbb{C}[\mathfrak{h}]^W \subset Q_c \subset \mathbb{C}[\mathfrak{h}]$. We prove that the algebra $\mathcal{D}(Q_c)$ of differential operators on quasi-invariants is a simple algebra, Morita equivalent to $\mathcal{D}(\mathfrak{h})$. The subalgebra $\mathcal{D}(Q_c)^W \subset \mathcal{D}(Q_c)$ of W -invariant operators turns out to be isomorphic to the spherical subalgebra $eH_{c,e} \subset H_c$. We show that $\mathcal{D}(Q_c)$ is generated, as an algebra, by Q_c and its ‘Fourier dual’ Q_c^b , and that $\mathcal{D}(Q_c)$ is a rank one projective $Q_c \otimes Q_c^b$ -module (via multiplication-action on $\mathcal{D}(Q_c)$ on opposite sides).

Table of Contents

1. Introduction
2. Standard modules over the rational Cherednik algebra
3. Harish-Chandra H_c -bimodules
4. The spherical subalgebra $eH_{c,e}$
5. A trace on the Cherednik algebra
6. The $eH_{c,e}$ -module structure on quasi-invariants
7. Differential operators on quasi-invariants
8. Translation functors and Morita equivalence
9. Applications of the shift operator
10. Appendix: A filtration on differential operators

1 Introduction

Let W be a finite Coxeter group in a complex vector space $\mathfrak{h}$, and $R \subset \mathfrak{h}^*$ the corresponding set of roots. To each W -invariant function $c : R \rightarrow \mathbb{C}$, $c \mapsto c_\alpha$, one can attach an associative algebra H_c , called the *rational Cherednik algebra*. This is a very interesting algebra tied to exciting works in combinatorics, completely integrable systems, and generalized McKay correspondence. Historically, the rational Cherednik algebra appeared as a ‘rational’ degeneration of the *double-affine Hecke algebra* introduced by Cherednik [Ch2]. Thus, the latter may (and should) be thought of as a deformation of the former. From this point of view, representation theory of the rational Cherednik algebra is perhaps ‘more basic’ than (or at least should be studied before) that of the double-affine Hecke algebra in the same sense as the representation theory of semisimple Lie algebras is ‘more basic’ than that of the corresponding quantum groups. Thus, one of our goals is to begin a systematic study of H_c representation theory.

Given $\alpha \in \mathfrak{h}^*$, write $\alpha^\vee \in \mathfrak{h}$ for the coroot, and $s_\alpha \in \text{GL}(\mathfrak{h})$ for the reflection corresponding to α . Recall from [EG], that the rational Cherednik algebra H_c (which was denoted $H_{1,c}$ in [EG]) is generated by the vector spaces $\mathfrak{h}$, $\mathfrak{h}^*$, and the set W , with defining relations (cf. formula (1.15) of [EG] for $t = 1$) given by

$$\begin{aligned}
 w \cdot x \cdot w^{-1} &= w(x) \quad , \quad w \cdot y \cdot w^{-1} = w(y) \quad , & \forall y \in \mathfrak{h} \quad , \quad x \in \mathfrak{h}^* \quad , \quad w \in W \\
 x_1 \cdot x_2 &= x_2 \cdot x_1 \quad , \quad y_1 \cdot y_2 = y_2 \cdot y_1 \quad , & \forall y_1, y_2 \in \mathfrak{h} \quad , \quad x_1, x_2 \in \mathfrak{h}^* \\
 y \cdot x - x \cdot y &= \langle y, x \rangle - \sum_{\alpha \in R/\{\pm 1\}} c_\alpha \cdot \langle y, \alpha \rangle \langle \alpha^\vee, x \rangle \cdot s_\alpha \quad , & \forall y \in \mathfrak{h} \quad , \quad x \in \mathfrak{h}^* .
 \end{aligned} \tag{1.1}$$

Thus, the elements $x \in \mathfrak{h}^*$ generate a subalgebra $\mathbb{C}[\mathfrak{h}] \subset H_c$ of polynomial functions on $\mathfrak{h}$, the elements $y \in \mathfrak{h}$ generate a subalgebra $\mathbb{C}[\mathfrak{h}^*] \subset H_c$, and the elements $w \in W$ span a copy of the group algebra $\mathbb{C}W$ sitting naturally inside H_c . Furthermore, it has been shown by Cherednik, see also [EG], that multiplication in H_c induces a vector space isomorphism:

$$\mathbb{C}[\mathfrak{h}] \otimes_{\mathbb{C}} \mathbb{C}W \otimes_{\mathbb{C}} \mathbb{C}[\mathfrak{h}^*] \xrightarrow{\sim} H_c \quad (\text{Poincaré-Birkhoff-Witt isomorphism for } H_c). \quad (1.2)$$

The name for the isomorphism above comes from its analogy with the well-known isomorphism: $U(\mathfrak{n}_-) \otimes_{\mathbb{C}} U(\mathfrak{h}) \otimes_{\mathbb{C}} U(\mathfrak{n}_+) \xrightarrow{\sim} U(\mathfrak{g})$ for the enveloping algebra of a complex semisimple Lie algebra $\mathfrak{g}$ with triangular decomposition: $\mathfrak{g} = \mathfrak{n}_+ + \mathfrak{h} + \mathfrak{n}_-$.

The Poincaré-Birkhoff-Witt isomorphism for H_c allows one to introduce a category $\mathcal{O}_{H_c}$ of modules over the algebra H_c similar to the category $\mathcal{O}_{\mathfrak{g}}$ of highest weight modules over $U(\mathfrak{g})$, considered by Bernstein-Gelfand-Gelfand, see [BGG]. The category $\mathcal{O}_{H_c}$ splits up into a direct sum of its subcategories $\mathcal{O}_{H_c}(\bar{\lambda})$, one for each $\bar{\lambda} \in \mathfrak{h}^*/W$. We will be mainly concerned below with the category $\mathcal{O}_{H_c}(0)$, which is most interesting among all $\mathcal{O}_{H_c}(\bar{\lambda})$'s. The category $\mathcal{O}_{H_c}(0)$ is the Cherednik algebra counterpart of the subcategory $\mathcal{O}_{\mathfrak{g}}(\chi) \subset \mathcal{O}_{\mathfrak{g}}$ corresponding, in the Bernstein-Gelfand-Gelfand setting¹, to a fixed character χ of the center of $U(\mathfrak{g})$.

The isomorphism (1.2) shows that the group algebra $\mathbb{C}W$ plays a role of the subalgebra $U(\mathfrak{h}) \subset U(\mathfrak{g})$. Thus, following the classical construction due to Verma, to each irreducible representation $\tau \in \text{Irrep}(W)$ one can associate a ‘standard’ module $M(\tau) \in \mathcal{O}_{H_c}(0)$, an analogue of Verma module. It is easy to show, see [DO], that each standard module $M(\tau)$ has a unique simple quotient, $L(\tau)$. Furthermore, any object of the category $\mathcal{O}_{H_c}(0)$ has finite length, and the collection $\{L(\tau)\}_{\tau \in \text{Irrep}(W)}$ is a complete collection of isomorphism classes of simple objects of $\mathcal{O}_{H_c}(0)$. Thus, the simple objects in $\mathcal{O}_{H_c}(0)$ are parametrized by the set $\text{Irrep}(W)$, while the simple objects in $\mathcal{O}_{\mathfrak{g}}(\chi)$ are parametrized (for regular χ) by elements of the Weyl group W itself.

The structure of the category $\mathcal{O}_{H_c}(0)$ depends crucially on the value of the parameter ‘ c ’. The category is semisimple for almost all ‘ c ’, see [OR], in which case $M(\tau) = L(\tau)$, for any τ . However, for a certain set of “singular” values of ‘ c ’, the multiplicities $[M(\tau) : L(\sigma)]$ are unknown. These multiplicities are Cherednik algebra analogues of Kazhdan-Lusztig type multiplicities for affine Hecke algebras, see [CG], and they are expected to be provided by some Intersection cohomology.

In this paper we are concerned with the case of *integral* values of ‘ c ’, which is, in a sense, intermediate between the two extreme cases above. We will see, although it is not a priori obvious², that, for any such ‘ c ’, the category $\mathcal{O}_{H_c}(0)$ is semisimple. Further, we apply H_c -representation theory to show that, for integral ‘ c ’, the algebra H_c is simple and Morita equivalent to $\mathcal{D}(\mathfrak{h}) \# W$, the cross product of W with the algebra $\mathcal{D}(\mathfrak{h})$ of polynomial differential operators on $\mathfrak{h}$.

Our motivation to study the case of integral ‘ c ’ comes also from an interesting connection with the theory of Calogero-Moser integrable systems and the theory of differential operators on singular algebraic varieties. In more detail, let $\mathbb{C}[\mathfrak{h}]^W \subset \mathbb{C}[\mathfrak{h}]$, be the subalgebra of W -invariant polynomials. For each non-negative integral value of ‘ c ’, Chalykh, Feigin, and Veselov [CV], [FV], have introduced an algebra Q_c of so-called *W-quasi-invariant* polynomials, such that $\mathbb{C}[\mathfrak{h}]^W \subset Q_c \subset \mathbb{C}[\mathfrak{h}]$. We use representation theory of Cherednik algebras to study differential operators on quasi-invariants, that is, the algebra $\mathcal{D}(Q_c)$ of differential operators on the singular variety $\text{Spec } Q_c$. The Calogero-Moser differential operator may be viewed as an analogue of the second order Laplacian on the variety $\text{Spec } Q_c$.

It turns out that the algebra $\mathcal{D}(Q_c)$ is almost as nice as the algebra of differential operators on a smooth variety. Specifically, we prove that the algebra $\mathcal{D}(Q_c)$ is Morita equivalent to the algebra

¹ We emphasize that it is the parameter ‘ c ’, and not $\bar{\lambda} \in \mathfrak{h}^*/W$ that plays the role of central character in representation theory of H_c . The parameter $\lambda \in \mathfrak{h}^*$ plays, in our present situation, the role of ‘Whittaker character’, see remark after Corollary 4.5.

² See remark at the end of §2, and also [OR].

of polynomial differential operators on the vector space $\mathfrak{h}$. The subalgebra $\mathcal{D}(Q_c)^W$ of W -invariant differential operators will be shown to be isomorphic to the spherical subalgebra $eH_c e$. Further, $\mathcal{D}(Q_c)$ is a simple algebra equipped with a natural involution, an analogue of the Fourier transform for differential operators on $\text{Spec } Q_c$. The algebra $\mathcal{D}(Q_c)$ contains Q_c and its ‘Fourier dual’, $Q_c^\flat$, as two maximal commutative subalgebras. We show that $\mathcal{D}(Q_c)$ is generated, as an algebra, by Q_c and $Q_c^\flat$, and that $\mathcal{D}(Q_c)$ is a rank one projective $Q_c \otimes Q_c^\flat$ -module (under multiplication-action on $\mathcal{D}(Q_c)$ on opposite sides).

The rings of differential operators on general singular algebraic varieties typically have rather unpleasant behavior (e.g. are not Noetherian, cf. [BGG1]). The question of simplicity and Morita equivalence of such rings has been studied by several authors, see [BW], [Sm], [Mu], [SS], [HS], [CS], [VdB], (and also [MvdB] for a result similar in spirit to ours). The varieties $\text{Spec } Q_c$ studied in the present paper seem to be natural generalizations of the one-dimensional examples constructed in [Sm] and [Mu].

The paper is organized as follows. Sections 2,3,5 are devoted to ‘pure’ representation theory of Cherednik algebras. In §2 we exploit an idea due to Opdam relating highest weight modules over the Cherednik algebra to finite dimensional representations of the Hecke algebra $H_W(e^{2\pi ic})$; thus §2 has some overlap with [DO] and [OR]. In section 3 a new and quite useful notion of a *Harish-Chandra H_c -bimodule* is introduced. In §4 we prove a Cherednik algebra counterpart of an important theorem (due to Levasseur-Stafford [LS]), saying that the spherical subalgebra in H_c is generated, for all regular values of ‘ c ’, by its two invariant commutative subalgebras, $\mathbb{C}[\mathfrak{h}]^W$ and $\mathbb{C}[\mathfrak{h}^*]^W$. This result has numerous applications. In §5 we compute a trace on the Cherednik algebra of type **A**, and use it to derive some applications to *finite-dimensional* representations of H_c . Quasi-invariants are introduced in §6, and the structure of the algebra $\mathcal{D}(Q_c)$ is studied in detail in §§7,9. The structure of $\mathcal{D}(Q_c)$ is in turn exploited in section 8 to get further results in representation theory of Cherednik algebras, e.g., to give an explicit construction of simple Harish-Chandra bimodules. These results of §8 bear some resemblance with the technique of *translation functors*, a well-known and very powerful tool in representation theory of semisimple Lie algebras.

Acknowledgments. We are very grateful to E. Opdam and R. Rouquier for generously sharing their ideas with us, and for making the results of [OR] available to us before its publication. These results play a crucial role in our arguments and, to a great extent, has triggered the present work. We would like to thank T. Stafford for many interesting comments and other useful information that was quite essential for us. The first author was partially supported by the NSF grant DMS 00-71792 and A. P. Sloan Research Fellowship; the work of the second author was partly conducted for the Clay Mathematics Institute and partially supported by the NSF grant DMS-9988796.

2 Standard modules over the rational Cherednik algebra

Most of the results of this section (in particular, Theorem 2.2 and Lemma 2.10) are due to Opdam-Rouquier [OR], and are reproduced here for the reader’s convenience only.

Fix a finite Coxeter group W in a complex vector space $\mathfrak{h}$. Thus, $\mathfrak{h}$ is the complexification of a real Euclidean vector space and W is generated by reflections with respect to a certain finite set $\{H_\alpha\}$ of hyperplanes in that Euclidean space. We write $(\cdot, \cdot)$ for the complex bilinear form on $\mathfrak{h}$ extending the Euclidean inner product. For each hyperplane, H_α , we choose nonzero linear functions $\pm\alpha \in \mathfrak{h}^*$ which vanish on H_α . The set $R \subset \mathfrak{h}^*$ of all such linear functions is called the set of roots. Write $s_\alpha \in \text{GL}(\mathfrak{h})$ for the reflection corresponding to $\alpha \in R$, and $\alpha^\vee \in \mathfrak{h}$ for the corresponding coroot, a vector such that $s_\alpha(\alpha^\vee) = -\alpha^\vee$. The lengths of roots and coroots are normalized so that $\langle \alpha, \alpha^\vee \rangle = 2$. We make a choice of the set R_+ of positive roots so that

$R = R_+ \sqcup (-R_+)$. The W -action on $\mathfrak{h}$ induces canonical actions on the symmetric algebras $\text{Sym}(\mathfrak{h}) = \mathbb{C}[\mathfrak{h}^*]$ and $\text{Sym}(\mathfrak{h}^*) = \mathbb{C}[\mathfrak{h}]$.

Let $\mathfrak{h}^{\text{reg}}$ denote the complement to the union of all the reflection hyperplanes, i.e., the complement to the zero set of the discriminant polynomial $\delta = \prod_{\alpha \in R_+} \alpha \in \mathbb{C}[\mathfrak{h}]$. Given a $\mathbb{C}[\mathfrak{h}]^W$ -module M , we will write $M|_{\mathfrak{h}^{\text{reg}}/W}$ for $\mathbb{C}[\mathfrak{h}^{\text{reg}}]^W \otimes_{\mathbb{C}[\mathfrak{h}]^W} M$, the localization to $\mathfrak{h}^{\text{reg}}/W$. Note that if M is a $\mathbb{C}[\mathfrak{h}]$ -module viewed as a $\mathbb{C}[\mathfrak{h}]^W$ -module by restriction of scalars, then $M|_{\mathfrak{h}^{\text{reg}}/W}$ is a $\mathbb{C}[\mathfrak{h}^{\text{reg}}]$ -module which coincides with $M|_{\mathfrak{h}^{\text{reg}}} := \mathbb{C}[\mathfrak{h}^{\text{reg}}] \otimes_{\mathbb{C}[\mathfrak{h}]} M$. In particular, $H_c|_{\mathfrak{h}^{\text{reg}}} = \mathbb{C}[\mathfrak{h}^{\text{reg}}] \otimes_{\mathbb{C}[\mathfrak{h}]} H_c$, the localization of the left regular H_c -module, acquires a natural algebra structure, hence an H_c -bimodule structure, such that the imbedding $H_c \hookrightarrow \mathbb{C}[\mathfrak{h}^{\text{reg}}] \otimes_{\mathbb{C}[\mathfrak{h}]} H_c$ becomes an algebra map. Alternatively, the algebra $H_c|_{\mathfrak{h}^{\text{reg}}}$ is obtained by Ore localization of H_c with respect to the multiplicative set $\{\delta^k\}_{k=1,2,\dots}$.

The group W acts freely on $\mathfrak{h}^{\text{reg}}$. Let B_W be the braid group of W , that is the fundamental group of the variety $\mathfrak{h}^{\text{reg}}/W$. Fix a point $*$ in $\mathfrak{h}^{\text{reg}}$ inside a Weyl chamber in $\mathfrak{h}$, and for each simple reflection $s_\alpha \in W$, let T_α be the class in $B_W = \pi_1(\mathfrak{h}^{\text{reg}}, *)$ corresponding to a straight path from the point $*$ to the point $s_\alpha(*)$ with an inserted little semi-circle (oriented counter-clockwise) around the hyperplane $\alpha = 0$. Given a W -invariant function $q : R \rightarrow \mathbb{C}^\times$, $\alpha \mapsto q_\alpha$, let $H_W(q)$ be the Hecke algebra. This algebra is obtained by taking the quotient of $\mathbb{C}[B_W]$ by the relations $(T_\alpha - 1)(T_\alpha + q_\alpha) = 0$, one for each simple reflection s_α . It is known that $\dim H_W(q) = |W|$, for any function $q : R \rightarrow \mathbb{C}^\times$.

Definition 2.1. A W -invariant function $q : R \rightarrow \mathbb{C}^\times$ is said to be *regular* if the Hecke algebra $H_W(q)$ is semisimple. Write Reg for the set of regular W -invariant functions q .

Since $H_W(1) = \mathbb{C}W$, we see that $q = 1$ is a regular function. Moreover, the set Reg is a dense Zariski open subset in the set of all W -invariant functions $q : R \rightarrow \mathbb{C}^\times$ (the latter set is naturally identified with $(\mathbb{C}^*)^l$, where l is the number of W -orbits in R). The complement of Reg has real codimension ≥ 2 , therefore Reg is a connected set. Hence, using rigidity of semisimple algebras one proves that $H_W(q) \simeq \mathbb{C}W$, for any $q \in \text{Reg}$.

We define standard modules over the Cherednik algebra H_c , see (1.1), as follows. Fix $\lambda \in \mathfrak{h}^*$, and let W_λ be the stabilizer of λ in W . Let $\mathbb{C}[\mathfrak{h}^*] \# W_\lambda$ be the cross-product of W_λ with the polynomial algebra. Sending $P \in \mathbb{C}[\mathfrak{h}^*]$ to $P(\lambda) \cdot 1$ yields an algebra homomorphism: $\mathbb{C}[\mathfrak{h}^*] \# W_\lambda \rightarrow \mathbb{C}W_\lambda$. Given $\tau \in \text{Irrep}(W_\lambda)$, an irreducible representation of W_λ , we write $\lambda \# \tau$ for the representation of $\mathbb{C}[\mathfrak{h}^*] \# W_\lambda$ obtained via the pull-back by the homomorphism above. Then we set $M(\lambda, \tau) := \text{Ind}_{\mathbb{C}[\mathfrak{h}^*] \# W_\lambda}^{H_c}(\lambda \# \tau)$. The module $M(\lambda, \tau)$ is called a *standard module*. In particular, if $\lambda = 0$ we have $W_\lambda = W$, in which case we write $M(0, \tau) = M(\tau)$ for the corresponding standard module.

Let $\mathbb{C}[R]^W$ denote the vector space of all W -invariant functions $c : R \rightarrow \mathbb{C}$, and $\mathbb{C}[R]_{\text{reg}}^W \subset \mathbb{C}[R]^W$ the set of functions $c \in \mathbb{C}[R]^W$ such that $\exp(2\pi ic) \in \text{Reg}$, i.e., such that the Hecke algebra $H_W(e^{2\pi ic})$ is semisimple. The goal of this section is to prove the following result due to Opdam-Rouquier in the key special case $\lambda = 0$.

Theorem 2.2 ([OR],[GGOR]). *If $c \in \mathbb{C}[R]_{\text{reg}}^W$ then, for any $\lambda \in \mathfrak{h}^*$ and $\tau \in \text{Irrep}(W_\lambda)$, the standard module $M(\lambda, \tau)$ is a simple H_c -module.*

Since $\exp(2\pi ic) = 1 \in \text{Reg}$ for any integral valued function c , Theorem 2.2 yields

Corollary 2.3. *All standard modules, $M(\lambda, \tau)$, are simple, for any $c \in \mathbb{Z}[R]^W$.* $\square$

By analogy with representation theory of semisimple Lie algebras we introduce the following

Definition 2.4. Let $\mathcal{O}_{H_c}$ be the category of finitely-generated H_c -modules M , such that the action on M of the subalgebra $\mathbb{C}[\mathfrak{h}^*] \subset H_c$ is locally finite, i.e., $\dim_{\mathbb{C}} \mathbb{C}[\mathfrak{h}^*] \cdot m < \infty$, for any $m \in M$.

We say that an object $M \in \mathcal{O}_{\mathbf{H}_c}$ has *type* $\bar{\lambda} \in \mathfrak{h}^*/W = \text{Spec}(\mathbb{C}[\mathfrak{h}^*]^W)$ if, for any $P \in \mathbb{C}[\mathfrak{h}^*]^W$, the action on M of the element $P - P(\bar{\lambda}) \in \mathbf{H}_c$ is locally nilpotent. Let $\mathcal{O}_{\mathbf{H}_c}(\bar{\lambda})$ be the full subcategory of modules having type $\bar{\lambda}$. Then, by a routine argument, cf. e.g. [Di], one obtains a direct sum decomposition: $\mathcal{O}_{\mathbf{H}_c} = \bigoplus_{\bar{\lambda} \in \mathfrak{h}^*/W} \mathcal{O}_{\mathbf{H}_c}(\bar{\lambda})$.

Lemma 2.5. (i) *Any object $M \in \mathcal{O}_{\mathbf{H}_c}$ is finitely generated over the subalgebra $\mathbb{C}[\mathfrak{h}] \subset \mathbf{H}_c$, in particular, $\mathcal{O}_{\mathbf{H}_c}$ is an abelian category.*

(ii) *For any $\lambda \in \mathfrak{h}^*$ and $\tau \in \text{Irrep}(W_\lambda)$, we have $M(\lambda, \tau) \in \mathcal{O}_{\mathbf{H}_c}$.*

(iii) *For any $M \in \mathcal{O}_{\mathbf{H}_c}$, there exists a nonzero homomorphism $M(\lambda, \tau) \rightarrow M$, for certain $\lambda \in \mathfrak{h}^*$ and $\tau \in \text{Irrep}(W_\lambda)$.*

(iv) *Every object of the category $\mathcal{O}_{\mathbf{H}_c}(0)$ has finite length.*

Remark. Using Lemma 2.5(iv), it is proven in [Gi] that, more generally, every object of the category $\mathcal{O}_{\mathbf{H}_c}$ also has finite length.

To prove Lemma 2.5, we need some notation. Let $\{x_i\}$ and $\{y_i\}$ be a pair of dual bases of $\mathfrak{h}^*$ and $\mathfrak{h}$, respectively. We view $\mathfrak{h}^*$ and $\mathfrak{h}$ as subspaces in $\mathbf{H}_c$, and let $\mathbf{h} = \frac{1}{2} \sum_i (x_i y_i + y_i x_i) \in \mathbf{H}_c$ denote the canonical element, which is independent of the choice of the bases. The element $\mathbf{h}$ satisfies the following commutation relations:

$$\mathbf{h} \cdot x = x \cdot (\mathbf{h} + 1), \quad \forall x \in \mathfrak{h}^*, \quad \text{and} \quad \mathbf{h} \cdot y = y \cdot (\mathbf{h} - 1), \quad \forall y \in \mathfrak{h}. \quad (2.6)$$

To prove the first of these formulas, use the notation $[a, b] := a \cdot b - b \cdot a$. Given $x \in \mathfrak{h}^*$, we calculate

$$\begin{aligned} [\mathbf{h}, x] &= \frac{1}{2} \left[\sum_i (x_i y_i + y_i x_i), x \right] = \frac{1}{2} \sum_i (x_i \cdot [y_i, x] + [y_i, x] \cdot x_i) \\ &= \frac{1}{2} \sum_i \left((x_i \cdot \langle y_i, x \rangle + \langle y_i, x \rangle \cdot x_i) - \sum_{\alpha \in R_+} \frac{c_\alpha}{2} (x_i \cdot \langle y_i, \alpha \rangle \langle \alpha^\vee, x \rangle s_\alpha + \langle y_i, \alpha \rangle \langle \alpha^\vee, x \rangle s_\alpha \cdot x_i) \right) \\ &= \sum_i (x_i \cdot \langle y_i, x \rangle) - \sum_{\alpha \in R_+} \frac{c_\alpha}{2} \cdot \langle \alpha^\vee, x \rangle \cdot \left(\sum_i (\langle y_i, \alpha \rangle x_i \cdot s_\alpha + \langle y_i, \alpha \rangle s_\alpha \cdot x_i) \right) \\ &= x - \sum_{\alpha \in R_+} \frac{c_\alpha}{2} \cdot \langle \alpha^\vee, x \rangle \cdot (\alpha \cdot s_\alpha + s_\alpha \cdot \alpha), \end{aligned} \quad (2.7)$$

where in the last equality we have used the identities $\sum_i \langle y_i, x \rangle \cdot x_i = x$, and $\sum_i \langle y_i, \alpha \rangle \cdot x_i = \alpha$. Since $(s_\alpha)^{-1} \cdot \alpha \cdot s_\alpha = s_\alpha(\alpha) = -\alpha$, we find

$$\alpha \cdot s_\alpha + s_\alpha \cdot \alpha = s_\alpha \cdot (s_\alpha)^{-1} \cdot \alpha \cdot s_\alpha + s_\alpha \cdot \alpha = s_\alpha \cdot (-\alpha) + s_\alpha \cdot \alpha = 0.$$

Thus, each term in the sum on the last line of (2.7) vanishes, and we deduce: $[\mathbf{h}, x] = x$. This proves the first identity in (2.6); the second one is proved similarly. $\square$

Let V be a bimodule over an associative algebra A . For any $a \in A$, we have an adjoint a -action on V given by $\text{ad } a : v \mapsto av - va$. We say that the adjoint action of A on V is *locally nilpotent* if, for any $v \in V$, there exists an integer $n \geq 0$ such that $\text{ad } a_0 \circ \text{ad } a_1 \circ \text{ad } a_2 \circ \dots \circ \text{ad } a_n(v) = 0$, for any $a_0, \dots, a_n \in A$. Let $n(v)$ be the smallest among such integers $n \geq 0$, to be referred to as the order of nilpotency of v .

Lemma 2.8. *Let A be a finitely generated commutative algebra and V an A -bimodule, such that the adjoint action of A on V is locally nilpotent. Then for any $v \in V$, the space AvA is finitely generated both as a left and as a right A -module.*

Proof. We proceed by induction in the order of nilpotency of v . If $n(v) = 0$, the result is clear, since $AvA = Av$. So, we have to prove the statement for $n(v) = m$, knowing it for $n(v) = m - 1$.

Fix $v \in V$ of nilpotency order m . Let $a_1, \dots, a_d$ be generators of A , and put: $u_i = [a_i, v]$, $i = 1, \dots, d$. We have: $A \cdot v \cdot A = \sum A \cdot v \cdot a_{i_1} \dots a_{i_N}$. By commuting v with a_{i_1} , we get $A \cdot v \cdot a_{i_1} \cdot a_{i_2} \dots a_{i_N} \subset A \cdot v \cdot a_{i_2} \dots a_{i_d} + A \cdot u_{i_1} \cdot A$. Now, continuing like this (i.e. interchanging v with a_{i_2} etc.), we get $A \cdot v \cdot A \subset A \cdot v + \sum_{i=1}^d A \cdot u_i \cdot A$. By the induction assumption, the module on the right is finitely generated. Thus, by the Hilbert-Noether lemma, so is the module AvA . $\square$

Proof of Lemma 2.5. (i) Let $M_0 \subset M$ be the finite dimensional vector space spanned by a finite set of generators. Then $H_c \cdot M_0 = M$. On the other hand, the space, $M'_0 := \mathbb{C}[\mathfrak{h}^*] \cdot M_0$ is finite dimensional, since the action of $\mathbb{C}[\mathfrak{h}^*]$ is locally finite. So $M = H_c \cdot M_0 = \mathbb{C}[\mathfrak{h}] \cdot CW \cdot \mathbb{C}[\mathfrak{h}^*] \cdot M_0 = \mathbb{C}[\mathfrak{h}] \cdot CW \cdot M'_0$. But $CW \cdot M'_0$ is finite dimensional, so we are done.

(ii) The H_c -module $M(\lambda, \tau)$ is generated by the W -stable finite dimensional subspace $E := CW \cdot \tau \subset M(\lambda, \tau)$. Hence, given $v \in M(\lambda, \tau)$, there exists a finite dimensional subspace $B \subset H_c$ such that $\mathbb{C}[\mathfrak{h}^*] \cdot v \subset \mathbb{C}[\mathfrak{h}^*] \cdot B \cdot E$. Moreover, since $\mathbb{C}[\mathfrak{h}^*]$ is finite over $\mathbb{C}[\mathfrak{h}^*]^W$, we may find B large enough, so that $\mathbb{C}[\mathfrak{h}^*] \cdot E \subseteq \mathbb{C}[\mathfrak{h}^*]^W \cdot B \cdot E$. But $\mathbb{C}[\mathfrak{h}^*]^W \cdot B \subset \mathbb{C}[\mathfrak{h}^*]^W \cdot B \cdot \mathbb{C}[\mathfrak{h}^*]^W$. By Lemma 2.8 applied to $V = H_c$, there exists a finite dimensional subspace $B' \subset H_c$, such that $\mathbb{C}[\mathfrak{h}^*]^W \cdot B \cdot \mathbb{C}[\mathfrak{h}^*]^W \subset B' \cdot \mathbb{C}[\mathfrak{h}^*]^W$. Hence, $\mathbb{C}[\mathfrak{h}^*]^W \cdot B \cdot E \subset \mathbb{C}[\mathfrak{h}^*]^W \cdot B \cdot \mathbb{C}[\mathfrak{h}^*]^W \cdot E \subset B' \cdot \mathbb{C}[\mathfrak{h}^*]^W \cdot E$. The latter space is finite dimensional since $\mathbb{C}[\mathfrak{h}^*]^W \cdot E$ is finite dimensional, and (ii) follows.

(iii) The action of the subalgebra $\mathbb{C}[\mathfrak{h}^*] \subset H_c$ on $M \in \mathcal{O}_{H_c}$ being locally finite, it follows that M contains a vector annihilated by the maximal ideal J of the algebra $\mathbb{C}[\mathfrak{h}^*]$ corresponding to a point $\lambda \in \mathfrak{h}^*$. The elements of M which are annihilated by J clearly form an W_λ -stable vector space. Hence, this vector space contains a simple W_λ -module σ . Any vector in this σ gives rise to a nonzero element of $\text{Hom}_{\mathbb{C}[\mathfrak{h}^*] \# W_\lambda}(\lambda \# \sigma, M) = \text{Hom}_{H_c}(M(\lambda, \sigma), M)$.

To prove (iv), we observe that the $\mathfrak{h}$ -action on any standard module $M(\tau)$ is diagonal, with finite dimensional eigenspaces. It follows that, for any $\tau \in \text{Irrep}(W)$, the multiplicity of each simple object of $\mathcal{O}_{H_c}(0)$ in $M(\tau)$ is bounded from above by the dimension of the corresponding $\mathfrak{h}$ -eigenspace. Further, the category $\mathcal{O}_{H_c}(0)$ has only finitely many simple objects, and therefore, each $M(\tau)$ has a finite Jordan-Hölder series.

We say that an object $N \in \mathcal{O}_{H_c}(0)$ is a highest weight object if it is a quotient of a standard module. It follows from the paragraph above that any highest weight object has finite length.

To prove that an arbitrary object $M \in \mathcal{O}_{H_c}(0)$ has finite length it suffices to show that M has a finite filtration by subobjects $0 = F_0 \subset F_1 \subset \dots \subset F_n = M$, such that successive quotients F_i/F_{i-1} are highest weight objects. To construct such a filtration, we use the fact that M contains a nonzero highest weight submodule N_1 , by part (iii) of the Lemma. Put $M_1 := M/N_1$, which is also an object of $\mathcal{O}_{H_c}(0)$. If $M = N_1$ we are done; if not, then M_1 also contains a nonzero highest weight object N_2 . We set $M_2 = M_1/N_2$, and iterate the procedure. Thus, for each $i \geq 1$, we get an object M_i which is a quotient of M . Let $F_i := \ker(M \twoheadrightarrow M_i)$. Clearly, $F_i \in \mathcal{O}_{H_c}(0)$, and we have an increasing chain of subobjects in M : $0 = F_0 \subset F_1 \subset \dots$. Since M is a Noetherian $\mathbb{C}[\mathfrak{h}]$ -module, this sequence must stabilize. This means that at some step l , the object N_{l+1} cannot be chosen, which means that $M_l = 0$. Thus, the $\{F_i\}$ provide a finite filtration with quotients being highest weight objects, and the result follows. $\square$

Remark. The argument above proves also that if for some $c \in \mathbb{C}[R]^W$ and $\bar{\lambda} \in \mathfrak{h}^*/W$, all the standard modules $M(\lambda, \tau)$ are simple, then every object of $\mathcal{O}_{H_c}(\bar{\lambda})$ has finite length. As we will see below, these conditions always hold, for instance, provided $c \in \mathbb{C}[R]_{\text{reg}}^W$.

Corollary. *The action of $\mathfrak{h}$ on any object $M \in \mathcal{O}_{H_c}(0)$ is locally finite, that is $\dim(\mathbb{C}[\mathfrak{h}] \cdot m) < \infty$, for any $m \in M$.* $\square$

We need to recall a few important results about Dunkl operators. According to Cherednik, see [EG] Proposition 4.5, the algebra H_c has a faithful "Dunkl representation", an injective algebra homomorphism $\Theta : H_c \hookrightarrow \mathcal{D}(\mathfrak{h}^{\text{reg}}) \# W$. This morphism extends by $\mathbb{C}[\mathfrak{h}^{\text{reg}}]$ -linearity to a map: $H_c|_{\mathfrak{h}^{\text{reg}}} = \mathbb{C}[\mathfrak{h}^{\text{reg}}] \otimes_{\mathbb{C}[\mathfrak{h}]} H_c \longrightarrow \mathcal{D}(\mathfrak{h}^{\text{reg}}) \# W$, which gives an algebra isomorphism $\Theta : H_c|_{\mathfrak{h}^{\text{reg}}} \xrightarrow{\sim} \mathcal{D}(\mathfrak{h}^{\text{reg}}) \# W$ (surjectivity is clear since the set of Dunkl operators generates $\mathcal{D}(\mathfrak{h}^{\text{reg}}) \# W$ over the subalgebra $\mathbb{C}[\mathfrak{h}^{\text{reg}}] \# W$). Now, for any H_c -module M , the localization, $M|_{\mathfrak{h}^{\text{reg}}}$, has a natural $H_c|_{\mathfrak{h}^{\text{reg}}}$ -module structure, therefore, a $\mathcal{D}(\mathfrak{h}^{\text{reg}}) \# W$ -module structure, via the Dunkl isomorphism above. Thus, since $\mathfrak{h}^{\text{reg}}$ is affine, one can view $M|_{\mathfrak{h}^{\text{reg}}}$ as a W -equivariant $\mathcal{D}$ -module on $\mathfrak{h}^{\text{reg}}$.

Observe further that the $\mathcal{D}$ -module on $\mathfrak{h}^{\text{reg}}$ arising from any object $M \in \mathcal{O}_{H_c}$ is finitely generated over the subalgebra $\mathbb{C}[\mathfrak{h}^{\text{reg}}]$, by Lemma 2.5(i). Hence, $M|_{\mathfrak{h}^{\text{reg}}}$, viewed as a $\mathcal{D}$ -module on $\mathfrak{h}^{\text{reg}}$, must be a vector bundle with flat connection. In particular, the standard module $M(\tau)$ is a free $\mathbb{C}[\mathfrak{h}]$ -module of rank $\dim(\tau)$, as follows from Poincaré-Birkhoff-Witt isomorphism (1.2). Thus $M(\tau)|_{\mathfrak{h}^{\text{reg}}}$, viewed as a $\mathcal{D}$ -module on $\mathfrak{h}^{\text{reg}}$, is the trivial vector bundle $\mathbb{C}[\mathfrak{h}^{\text{reg}}] \otimes \tau$ equipped with a flat connection.

The following well-known result plays a crucial role in this paper.

Proposition 2.9. (i) (Dunkl [D], Cherednik [Ch]) *The connection on $\mathbb{C}[\mathfrak{h}^{\text{reg}}] \otimes \tau$ arising from the standard module $M(\tau)$ is the Knizhnik-Zamolodchikov connection with values in τ .*

(ii) (Opdam [O]) *The monodromy representation of the fundamental group $B_W = \pi_1(\mathfrak{h}^{\text{reg}}/W)$ corresponding to this connection factors through the Hecke algebra $H_W(e^{2\pi ic})$.*

Proof. Although this is a known result, we will give a short proof for reader's convenience. Let us write down a system of linear differential equations which defines the horizontal sections of the corresponding connection. For this purpose, consider the generating subspace $\tau \subset M(\tau)|_{\mathfrak{h}^{\text{reg}}}$. We have $yv = 0$ for $y \in \mathfrak{h}$ and $v \in \tau$. Therefore (since the isomorphism $\Theta : H_c|_{\mathfrak{h}^{\text{reg}}} \rightarrow \mathcal{D}(\mathfrak{h}^{\text{reg}}) \# W$ is defined via Dunkl operators), we have

$$\partial_y v = \left[y - \sum_{\alpha \in R_+} c_\alpha \frac{\langle \alpha, y \rangle}{\alpha} (s_\alpha - 1) \right] v = \sum_{\alpha \in R_+} c_\alpha \frac{\langle \alpha, y \rangle}{\alpha} (1 - s_\alpha) v$$

In the trivialization $M(\tau)|_{\mathfrak{h}^{\text{reg}}} \simeq \mathbb{C}[\mathfrak{h}^{\text{reg}}] \otimes \tau$, the above formulas for the Dunkl operators equip the trivial bundle $\mathbb{C}[\mathfrak{h}^{\text{reg}}] \otimes \tau$ with the flat connection

$$\nabla = d + \sum_{\alpha \in R_+} c_\alpha \frac{d\alpha}{\alpha} \otimes (\text{Id}_\tau - s_\alpha) : \mathbb{C}[\mathfrak{h}^{\text{reg}}] \otimes \tau \longrightarrow \Omega^1(\mathfrak{h}^{\text{reg}}) \otimes \tau.$$

This vector bundle with connection ∇ is W -equivariant, hence, descends to $\mathfrak{h}^{\text{reg}}/W$. The corresponding flat sections are (multivalued) functions $f : \mathfrak{h}^{\text{reg}}/W \rightarrow \tau$ satisfying the holonomic system of differential equations: $\partial_y f = \sum_{\alpha \in R_+} c_\alpha \frac{\langle \alpha, y \rangle}{\alpha} (1 - s_\alpha) f$. An elementary theory of ordinary differential equations says that, for generic c , the image of any element T_α in the monodromy representation of this system satisfies the equation $(T_\alpha - 1)(T_\alpha + e^{2\pi ic_\alpha}) = 0$. By continuity, the same holds for all c , see [O] for more details. $\square$

In general, given a W -equivariant vector bundle $\mathcal{M}$ on $\mathfrak{h}^{\text{reg}}$ with flat connection, the germs of horizontal holomorphic sections of $\mathcal{M}$ form a locally constant sheaf on $\mathfrak{h}^{\text{reg}}/W$. Let $\text{Mon}(\mathcal{M})$ be the corresponding monodromy representation of the fundamental group $\pi_1(\mathfrak{h}^{\text{reg}}/W, *)$ in the fiber over $*$, where $*$ is some fixed point in $\mathfrak{h}^{\text{reg}}/W$. The assignment: $\mathcal{M} \mapsto \text{Mon}(\mathcal{M})$ gives a functor from the category of W -equivariant vector bundles on $\mathfrak{h}^{\text{reg}}$ with flat connection to the category of finite dimensional representations of the group $\pi_1(\mathfrak{h}^{\text{reg}}/W, *) = B_W$.

The following result is due to Opdam-Rouquier.

Lemma 2.10 ([OR],[GGOR]). *Let N be an object of $\mathcal{O}_{\mathfrak{H}_c}$ which is torsion-free over the subalgebra $\mathbb{C}[\mathfrak{h}] \subset \mathfrak{H}_c$. Then the canonical map: $\mathrm{Hom}_{\mathfrak{H}_c}(M, N) \rightarrow \mathrm{Hom}_{B_W}(\mathrm{Mon}(M|_{\mathfrak{h}^{\mathrm{reg}}}), \mathrm{Mon}(N|_{\mathfrak{h}^{\mathrm{reg}}}))$ is injective, for any $M \in \mathcal{O}_{\mathfrak{H}_c}$.*

Proof. (borrowed from [OR],[GGOR]). By assumption, the module N contains no $\mathbb{C}[\mathfrak{h}]$ -submodule supported on a proper subset of $\mathfrak{h}$. It follows that if $f : M \rightarrow N$ is an $\mathfrak{H}_c$ -module morphism such that the induced morphism: $M|_{\mathfrak{h}^{\mathrm{reg}}} \rightarrow N|_{\mathfrak{h}^{\mathrm{reg}}}$ has zero image, then f itself has the zero image, that is $f = 0$. Thus, for any $M \in \mathcal{O}_{\mathfrak{H}_c}$, the canonical map $i_1 : \mathrm{Hom}_{\mathfrak{H}_c}(M, N) \rightarrow \mathrm{Hom}_{\mathfrak{H}_c|_{\mathfrak{h}^{\mathrm{reg}}}}(M|_{\mathfrak{h}^{\mathrm{reg}}}, N|_{\mathfrak{h}^{\mathrm{reg}}})$ is injective.

Now, as has been explained earlier, we may regard $\mathcal{M} := M|_{\mathfrak{h}^{\mathrm{reg}}}$ and $\mathcal{N} := N|_{\mathfrak{h}^{\mathrm{reg}}}$ as W -equivariant vector bundles on $\mathfrak{h}^{\mathrm{reg}}$ with flat connections. Hence, the map of the Lemma can be factored as a composition

$$\begin{aligned} \mathrm{Hom}_{\mathfrak{H}_c}(M, N) &\xrightarrow{i_1} \mathrm{Hom}_{\mathfrak{H}_c|_{\mathfrak{h}^{\mathrm{reg}}}}(M|_{\mathfrak{h}^{\mathrm{reg}}}, N|_{\mathfrak{h}^{\mathrm{reg}}}) \\ &= \mathrm{Hom}_{\mathcal{D}(\mathfrak{h}^{\mathrm{reg}})\#W}(\mathcal{M}, \mathcal{N}) \xrightarrow{\sim} \left\{ \begin{array}{c} \text{global } W\text{-invariant horizontal} \\ \text{sections of } (\mathcal{M}^* \otimes \mathcal{N}) \end{array} \right\} \\ &\xrightarrow{i_2} (\mathrm{Mon}(\mathcal{M}^* \otimes \mathcal{N}))^{B_W} = \mathrm{Hom}_{B_W}(\mathrm{Mon}(M|_{\mathfrak{h}^{\mathrm{reg}}}), \mathrm{Mon}(N|_{\mathfrak{h}^{\mathrm{reg}}})). \end{aligned}$$

We already know that i_1 is injective. Further, any $\mathcal{D}(\mathfrak{h}^{\mathrm{reg}})\#W$ -module morphism between vector bundles with flat connections is given by a horizontal section of the Hom-bundle. The latter is nothing but a global horizontal section of $\mathcal{M}^* \otimes \mathcal{N}$, hence i_2 is injective. Finally, assigning to a horizontal section its value in the fiber at a given point $* \in \mathfrak{h}^{\mathrm{reg}}/W$ gives an injection (which is a bijection, if the connection has regular singularities), and the Lemma follows. $\square$

The following corollary of the above results and its proof were communicated to us by E. Opdam.

Corollary 2.11 ([OR],[GGOR]). *If $c \in \mathbb{C}[R]_{\mathrm{reg}}^W$, then all standard modules $M(\tau)$ are irreducible.*

Proof. Fix $\tau \in \mathrm{Irrep}(W)$, and given $c \in \mathbb{C}[R]_{\mathrm{reg}}^W$ write $\mathrm{Mon}_{\tau}(c) := \mathrm{Mon}(M(\tau)|_{\mathfrak{h}^{\mathrm{reg}}})$ for the corresponding monodromy representation of the Hecke algebra $H_W(e^{2\pi ic})$. The assignment $c \mapsto \mathrm{Mon}_{\tau}(c)$ gives a continuous function on the set $\mathbb{C}[R]_{\mathrm{reg}}^W$. Furthermore, this set is connected and contains the point $c = 0$, for which we have: $\mathrm{Mon}_{\tau}(0) = \tau$. Since, $H_W(q) \simeq \mathbb{C}W$ for all $q \in \mathrm{Reg}$, a simple deformation argument shows that $\mathrm{Mon}_{\tau}(c)$ is a simple $H_W(e^{2\pi ic})$ -module, for any $c \in \mathbb{C}[R]_{\mathrm{reg}}^W$. Since $M(\tau)$ is free over $\mathbb{C}[\mathfrak{h}]$, hence torsion-free, Lemma 2.10 yields: $\dim \mathrm{Hom}_{\mathfrak{H}_c}(M(\tau), M(\tau)) = 1$.

Next, fix two non-isomorphic W -modules $\tau, \sigma \in \mathrm{Irrep}(W)$. A similar argument shows that, for each $c \in \mathbb{C}[R]_{\mathrm{reg}}^W$, the $H_W(e^{2\pi ic})$ -modules $\mathrm{Mon}_{\tau}(c)$ and $\mathrm{Mon}_{\sigma}(c)$ are simple and non-isomorphic to each other. Thus, $\mathrm{Hom}_{B_W}(\mathrm{Mon}_{\sigma}(c), \mathrm{Mon}_{\tau}(c)) = 0$. Hence, Lemma 2.10 yields: $\mathrm{Hom}_{\mathfrak{H}_c}(M(\sigma), M(\tau)) = 0$. Thus, we conclude: $\dim \mathrm{Hom}_{\mathfrak{H}_c}(M(\sigma), M(\tau)) = \dim \mathrm{Hom}_W(\tau, \sigma)$, $\forall \sigma, \tau \in \mathrm{Irrep}(W)$.

Now, assume $M(\tau)$ is not a simple $\mathfrak{H}_c$ -module, and let $M \subset M(\tau)$ be a proper nonzero submodule. Then, Lemma 2.5(iii) says that there exists $\lambda \in \mathfrak{h}^*$ and $\sigma \in \mathrm{Irrep}(W_{\lambda})$ such that $\mathrm{Hom}_{\mathfrak{H}_c}(M(\lambda, \sigma), M) \neq 0$. Furthermore, the direct sum decomposition $\mathcal{O}_{\mathfrak{H}_c} = \bigoplus_{\bar{\lambda} \in \mathfrak{h}^*/W} \mathcal{O}_{\mathfrak{H}_c}(\bar{\lambda})$ implies that λ must be zero, so that $M(\lambda, \sigma) = M(\sigma)$ and $\sigma \in \mathrm{Irrep}(W)$. Hence, the dimension of $\mathrm{Hom}_{\mathfrak{H}_c}(M(\sigma), M(\tau))$ is at least 1 if $\sigma \neq \tau$, and is at least 2 if $\sigma = \tau$. But this contradicts the dimension equality: $\dim \mathrm{Hom}_{\mathfrak{H}_c}(M(\sigma), M(\tau)) = \dim \mathrm{Hom}_W(\tau, \sigma)$ proved earlier. The contradiction shows that $M(\tau)$ is simple. $\square$

The module $M(\lambda, \tau)$ can be identified, as a vector space, with $\mathbb{C}[\mathfrak{h}] \otimes \text{Ind}_{W_\lambda}^W \tau$, via the PBW-theorem. We fix such an identification. Thus, the standard increasing filtration on $\mathbb{C}[\mathfrak{h}]$ by degree of polynomials gives rise to an increasing filtration $F_\bullet M(\lambda, \tau)$ on $M(\lambda, \tau)$, such that $F_0 M(\lambda, \tau) = \text{Ind}_{W_\lambda}^W \tau$. It is clear that operators corresponding to elements $y \in \mathfrak{h}$ preserve this filtration.

Lemma 2.12. *If $c \in \mathbb{C}[R]_{\text{reg}}^W$, then any $\mathfrak{h}$ -weight vector in $M(\lambda, \tau)$ belongs to $F_0 M(\lambda, \tau)$.*

Proof. The proof is based on a continuity argument and on the fact that the result clearly holds true, provided $M(\lambda, \tau)$ is a simple $\mathbf{H}_c$ -module, e.g., for $\lambda = 0$.

First, we form the induced module $X(\lambda) = \text{Ind}_{\mathbb{C}[\mathfrak{h}^*]}^{\mathbf{H}_c} \lambda$. We can (and will) canonically identify, via the PBW-theorem, the underlying vector space of the module $X(\lambda)$ with $\mathbb{C}[\mathfrak{h}] \otimes \mathbb{C}W$. As above, the standard increasing filtration on $\mathbb{C}[\mathfrak{h}]$ by degree of polynomials gives rise to an increasing filtration $F_\bullet X(\lambda)$, such that $F_0 X(\lambda) = \mathbb{C}W$.

Let $\text{Weight } X(\lambda)$ be the sum of all the $\mathfrak{h}$ -weight subspaces of $X(\lambda)$. It is clear that, for any $\lambda \in \mathfrak{h}^*$, the space $\text{Weight } X(\lambda)$ contains $F_0 X(\lambda)$. Further, one has an $\mathbf{H}_c$ -module direct sum decomposition: $X(\lambda) = \bigoplus_{\tau \in \text{Irrep}(W_\lambda)} \tau^* \otimes M(\lambda, \tau)$, where τ^* stands for the representation dual to τ . Therefore, to prove the Lemma it suffices to show that every $\mathfrak{h}$ -weight vector in $X(\lambda)$ belongs to $F_0 X(\lambda) = \mathbb{C}W$.

Fix $\lambda \in \mathfrak{h}^*$, and for any $t \in \mathbb{C}$, view $\text{Weight } X(t \cdot \lambda)$ as a subspace in $\mathbb{C}[\mathfrak{h}] \otimes \mathbb{C}W$ (note that the latter is *independent* of $t \in \mathbb{C}$). For any integer $d \geq 0$, consider the set

$$U_d = \{t \in \mathbb{C} \mid (\text{Weight } X(t \cdot \lambda)) \cap F_d X(t \cdot \lambda) = F_0 X(t \cdot \lambda)\}.$$

It is clear that $\mathbb{C} = U_0 \supset U_1 \supset \dots$. Observe that by Corollary 2.11, $M(\tau)$ is a simple $\mathbf{H}_c$ -module, for any $\tau \in \text{Irrep}(W)$. It follows that for $t = 0$ one has $\text{Weight } X(t \cdot \lambda) = F_0 X(t \cdot \lambda)$. Hence, each set U_d contains the point $t = 0$ and is Zariski open, by an elementary continuity argument. Hence, for each $d \geq 0$, the set $\mathbb{C} \setminus U_d$ is a proper Zariski closed, hence finite, subset of $\mathbb{C}$. It follows that the set $\bigcup_{d \geq 0} (\mathbb{C} \setminus U_d)$ is a union of a countable family of finite sets, hence a countable subset in $\mathbb{C}$.

Let T be the set of all $t \in \mathbb{C}$ such that $\text{Weight } X(t \cdot \lambda) = F_0 X(t \cdot \lambda)$. By definition we have $T = \bigcap_{d \geq 0} U_d$. Therefore, $\mathbb{C} \setminus T = \bigcup_{d \geq 0} (\mathbb{C} \setminus U_d)$ is a countable set. Further, this countable set must be invariant under dilations, since the assignment $x \mapsto t \cdot x$, $y \mapsto t^{-1} \cdot y$, $t \in \mathbb{C}^*$, $x \in \mathfrak{h}^*$, $y \in \mathfrak{h}$, gives a $\mathbb{C}^*$ -action on $\mathbf{H}_c$ by algebra automorphisms. Finally, we know that $0 \in T$, i.e., $0 \notin \mathbb{C} \setminus T$. These properties force the set $\mathbb{C} \setminus T$ to be empty, hence $T = \mathbb{C}$, and we are done. $\square$

Proof of Theorem 2.2. Let N be a nonzero submodule of $M(\lambda, \tau)$. Thus, $N \in \mathcal{O}_{\mathbf{H}_c}$. Hence, N contains a nonzero $\mathfrak{h}$ -weight vector v . Lemma 2.12 yields $v \in F_0 M(\lambda, \tau)$. But $F_0 M(\lambda, \tau) = \text{Ind}_{W_\lambda}^W \tau$ is an irreducible $\mathbb{C}[\mathfrak{h}^*] \# W$ -module, so $v \in F_0 M(\lambda, \tau) \subset N$ implies $N = M(\lambda, \tau)$. $\square$

Remark. It has been shown in [DO] that each standard module $M(\tau)$, $\tau \in \text{Irrep}(W)$, has a unique simple quotient $L(\tau)$. Moreover, any object of the category $\mathcal{O}_{\mathbf{H}_c}(0)$ has finite length, and the finite set $\{L(\tau)\}_{\tau \in \text{Irrep}(W)}$ is a complete collection of the isomorphism classes of simple objects of $\mathcal{O}_{\mathbf{H}_c}(0)$. Further, it is proved in [Gu] and [OR] that the category $\mathcal{O}_{\mathbf{H}_c}(0)$ has enough projectives, i.e., each module $L(\tau)$ has an indecomposable projective cover $P(\tau) \in \mathcal{O}_{\mathbf{H}_c}(0)$. Moreover, any projective $P \in \mathcal{O}_{\mathbf{H}_c}(0)$ has a finite increasing filtration by $\mathbf{H}_c$ -submodules: $0 = F_0 \subset F_1 \subset \dots \subset F_l = P$, such that $F_i/F_{i-1} \simeq M(\sigma_i)$, for some standard modules $M(\sigma_i)$, $\sigma_i \in \text{Irrep}(W)$, $i = 1, \dots, l$. Furthermore, for any $\tau, \sigma \in \text{Irrep}(W)$, Guay [Gu], and Opdam-Rouquier [OR], proved the following analogue of Brauer-Bernstein-Gelfand-Gelfand type reciprocity formula $[P(\tau) : M(\sigma)] = [M(\sigma) : L(\tau)]$, see [BGG]. This formula implies, in particular, that the category $\mathcal{O}_{\mathbf{H}_c}(0)$ is semisimple if and only if all the standard modules $M(\tau)$, $\tau \in \text{Irrep}(W)$, are simple, i.e. if and only if $M(\tau) = L(\tau)$, $\forall \tau \in \text{Irrep}(W)$. It would be interesting to obtain similar results for the category $\mathcal{O}_{\mathbf{H}_c}(\bar{\lambda})$, with $\bar{\lambda} \neq 0$.

3 Harish-Chandra H_c -bimodules

One of the goals of this section is to prove

Theorem 3.1. *If $c \in \mathbb{C}[R]_{\text{reg}}^W$, then H_c is a simple algebra.*

The proof of the Theorem will be based on the concept of a Harish-Chandra bimodule which we now introduce.

Let V be an H_c -bimodule, i.e., a left $H_c \otimes H_c^{\text{op}}$ -module. For any $x \in H_c$ we have an adjoint x -action on V given by $\text{ad } x : v \mapsto xv - vx$.

Definition 3.2. A finitely generated $H_c \otimes (H_c)^{\text{op}}$ -module V is called a Harish-Chandra bimodule if, for any $x \in \mathbb{C}[\mathfrak{h}]^W$ or $x \in \mathbb{C}[\mathfrak{h}^*]^W$, the $\text{ad } x$ -action on V is locally nilpotent.

Here are the first elementary results about Harish-Chandra bimodules.

Lemma 3.3. (i) *Harish-Chandra bimodules form a full abelian subcategory in the category of H_c -bimodules; this subcategory is stable under extensions.*

(ii) *Any Harish-Chandra bimodule is finitely-generated over the subalgebra $\mathbb{C}[\mathfrak{h}]^W \otimes \mathbb{C}[\mathfrak{h}^*]^W \subset H_c \otimes (H_c)^{\text{op}}$, with $\mathbb{C}[\mathfrak{h}^*]^W$ acting on the right and $\mathbb{C}[\mathfrak{h}]^W$ on the left. Similarly, it is also finitely-generated over the subalgebra $\mathbb{C}[\mathfrak{h}^*]^W \otimes \mathbb{C}[\mathfrak{h}]^W \subset H_c \otimes (H_c)^{\text{op}}$;*

(iii) *Any Harish-Chandra bimodule is finitely-generated as a left, resp. right, H_c - module;*

(iv) *For any maximal ideal $J \subset \mathbb{C}[\mathfrak{h}^*]$ and any Harish-Chandra bimodule V , we have $V/V \cdot J \in \mathcal{O}_{H_c}$, as a left module.*

(v) *The algebra H_c is a Harish-Chandra H_c -bimodule.*

Proof. The algebra H_c has an increasing filtration $F_\bullet H_c$ such that $\text{gr}(H_c) = \mathbb{C}[\mathfrak{h} \times \mathfrak{h}^*] \# W$, see [EG]. Hence H_c is a Noetherian algebra, for this is clearly true for the algebra $\text{gr}(H_c) = \mathbb{C}[\mathfrak{h} \times \mathfrak{h}^*] \# W$. Part (i) of the Lemma follows.

To prove (ii), observe that the filtration on H_c gives rise to the tensor product filtration: $(H_c \otimes H_c^{\text{op}})_p := \sum_{p=i+j} F_i(H_c) \otimes F_j(H_c^{\text{op}})$, on the algebra $H_c \otimes H_c^{\text{op}}$. Choose a finite-dimensional subspace $V_0 \subset V$ generating V as an $H_c \otimes H_c^{\text{op}}$ -module and, for each $p \geq 0$, put $V_p := (H_c \otimes H_c^{\text{op}})_p \cdot V_0$. This is a good filtration on V , and one can regard $\text{gr}(V)$, the associated graded space, as a $W \times W$ -equivariant finitely-generated module over $\mathbb{C}[\mathfrak{h}_1 \times \mathfrak{h}_2 \times \mathfrak{h}_1^* \times \mathfrak{h}_2^*]$, where the subscripts ‘1, 2’ indicate the corresponding copy of $\mathfrak{h}$. Then, for any homogeneous element $P \in \mathbb{C}[\mathfrak{h}]^W$, the $\text{ad } P$ -action on V corresponds to the action of the element $P^{\text{left}} \otimes 1 - 1 \otimes P^{\text{right}} \in \mathbb{C}[\mathfrak{h}_1]^W \otimes \mathbb{C}[\mathfrak{h}_2]^W$ on $\text{gr}(V)$. The (local) nilpotency of the $\text{ad } P$ -action on a Harish-Chandra module thus implies that, if P is homogeneous of degree > 0 , then the action of the element $P^{\text{left}} \otimes 1 - 1 \otimes P^{\text{right}}$ on $\text{gr}V$ is nilpotent. Hence, the support of $\text{gr}V$, viewed as a $\mathbb{C}[\mathfrak{h}_1 \times \mathfrak{h}_2 \times \mathfrak{h}_1^* \times \mathfrak{h}_2^*]$ -module, is contained in the zero locus of the polynomial $P^{\text{left}} \otimes 1 - 1 \otimes P^{\text{right}} \in \mathbb{C}[\mathfrak{h}_1 \times \mathfrak{h}_2 \times \mathfrak{h}_1^* \times \mathfrak{h}_2^*]$. This way, for any Harish-Chandra module V , we deduce the following upper bound on the set $\text{supp}(\text{gr}V) \subset \mathfrak{h}_1 \times \mathfrak{h}_2 \times \mathfrak{h}_1^* \times \mathfrak{h}_2^*$:

$$\text{supp}(\text{gr}V) \subset \left(\bigcup_{w \in W} \text{Graph}(w) \right) \times \left(\bigcup_{y \in W} \text{Graph}(y) \right), \quad (3.4)$$

where $\text{Graph}(w)$ denotes the graph in $\mathfrak{h}_1 \times \mathfrak{h}_2$, resp. $\mathfrak{h}_1^* \times \mathfrak{h}_2^*$, of the w -action map: $x \mapsto w(x)$.

In particular, the restriction of the composite map: $\mathfrak{h}_1 \times \mathfrak{h}_2 \times \mathfrak{h}_1^* \times \mathfrak{h}_2^* \rightarrow \mathfrak{h}_1 \times \mathfrak{h}_1^* \rightarrow \mathfrak{h}_1/W \times \mathfrak{h}_1^*/W$ to $\text{supp}(\text{gr}V)$ is a finite map. Therefore, $\text{gr}(V)$ is finitely generated over the subalgebra $\mathbb{C}[\mathfrak{h}]^W \otimes \mathbb{C}[\mathfrak{h}^*]^W$, hence the same holds for V itself. Part (ii) follows.

To prove (iii) we observe that (3.4) implies also that the restriction to $\text{supp}(\text{gr}V)$ of the corresponding projection: $\mathfrak{h}_1 \times \mathfrak{h}_2 \times \mathfrak{h}_1^* \times \mathfrak{h}_2^* \rightarrow \mathfrak{h}_1 \times \mathfrak{h}_2^*$ is a finite map. We deduce similarly that

$\mathfrak{gr}V$ is finitely generated with respect to the left action of the algebra $\mathfrak{gr}(\mathbf{H}_c)$. This forces V to be finitely generated left $\mathbf{H}_c$ -module, and (iii) follows.

Clearly (ii) implies (iv). To prove (v), endow $\mathbf{H}_c$ with an increasing filtration by assigning elements of $\mathfrak{h}$ filtration degree 1, and elements of $\mathfrak{h}^*$ and $\mathbb{C}W$ filtration degree 0 (this is not the standard filtration $F_\bullet \mathbf{H}_c$ used earlier). The defining relations (1.1) of the Cherednik algebra show that commuting with an element of $\mathbb{C}[\mathfrak{h}^*]^W \subset \mathbf{H}_c$ decreases filtration degree at least by 1. Hence, the adjoint action of $\mathbb{C}[\mathfrak{h}^*]^W$ is locally nilpotent. The other condition is proved similarly. $\square$

Proposition 3.5. *If $c \in \mathbb{C}[R]_{\text{reg}}^W$, then any Harish-Chandra $\mathbf{H}_c$ -bimodule is a finite rank projective module over the algebra $\mathbb{C}[\mathfrak{h}]^W \otimes \mathbb{C}[\mathfrak{h}^*]^W \subset \mathbf{H}_c \otimes \mathbf{H}_c^{\text{op}}$.*

Proof. Recall that a finitely generated module F over A , a commutative finitely generated algebra without zero-divisors, is projective if and only if, for all maximal ideals $\mathfrak{m} \subset A$, the geometric fibers $F/\mathfrak{m} \cdot F$ all have the same (finite) dimension over $\mathbb{C}$. Thus, we arrive at the following

Observation: *Let X, Y be irreducible algebraic varieties, and $\mathcal{F}$ a coherent sheaf on $X \times Y$. Suppose that for any $x \in X$, the sheaf $\mathcal{F}_x = \mathcal{F}|_{x \times Y}$ is locally free of finite rank, and for any $y \in Y$ the sheaf $\mathcal{F}_y = \mathcal{F}|_{X \times y}$ is locally free of finite rank. Then $\mathcal{F}$ is itself locally free of finite rank.*

Now let V be a Harish-Chandra $\mathbf{H}_c \otimes \mathbf{H}_c^{\text{op}}$ -module. We view V as a module over $\mathbb{C}[\mathfrak{h}]^W \otimes \mathbb{C}[\mathfrak{h}^*]^W$, with $\mathbb{C}[\mathfrak{h}^*]$ acting on the right and $\mathbb{C}[\mathfrak{h}]$ on the left. By the previous Lemma, this module is finitely generated. Therefore, by the Observation above, to prove the Proposition we only have to show that, for any maximal ideal $J \subset \mathbb{C}[\mathfrak{h}^*]^W$, the left $\mathbb{C}[\mathfrak{h}]^W$ -module $V/V \cdot J$ is locally free of finite rank (and a similar result for a right $\mathbb{C}[\mathfrak{h}^*]^W$ -module $V/I \cdot V$, $I \subset \mathbb{C}[\mathfrak{h}]^W$). Thus, let $J \subset \mathbb{C}[\mathfrak{h}^*]^W$ be the maximal ideal, corresponding to a point $\bar{\lambda} \in \mathfrak{h}^*/W$. Then $V/V \cdot J \in \mathcal{O}_{\mathbf{H}_c}(\bar{\lambda})$. This module has a finite composition series whose successive quotients are standard modules, since $c \in \mathbb{C}[R]_{\text{reg}}^W$. But all standard modules, $M(\lambda, \tau)$, are $\mathbb{C}[\mathfrak{h}]^W$ -free, and the statement follows. $\square$

Let $V|_{\mathfrak{h}^{\text{reg}}}$ denote the localization of a Harish-Chandra $\mathbf{H}_c$ -bimodule V with respect to the *left* action of the subalgebra $\mathbb{C}[\mathfrak{h}] \subset \mathbf{H}_c$. We note that $V|_{\mathfrak{h}^{\text{reg}}}$ coincides also with the localization of V with respect to the *right* action of the subalgebra $\mathbb{C}[\mathfrak{h}]$. To see this, observe that in either case, one may replace localizing with respect to $\mathbb{C}[\mathfrak{h}]$ by localizing with respect to the smaller algebra $\mathbb{C}[\mathfrak{h}]^W$. But it follows easily from Definition 3.2 that left and right localizations of V with respect to $\mathbb{C}[\mathfrak{h}]^W$ coincide.

Proposition 3.5 yields readily

Corollary 3.6. *Let $c \in \mathbb{C}[R]_{\text{reg}}^W$, and let V be a Harish-Chandra $\mathbf{H}_c$ -bimodule such that $V|_{\mathfrak{h}^{\text{reg}}} = 0$. Then $V = 0$.* $\square$

Proof of Theorem 3.1. Let $I \subset \mathbf{H}_c$ be a nonzero two-sided ideal. Then $V := \mathbf{H}_c/I$ is, by Lemma 3.3(v), a Harish-Chandra bimodule. Moreover, since $\mathbf{H}_c$ is a free left $\mathbb{C}[\mathfrak{h}]$ -module, and any submodule of a free $\mathbb{C}[\mathfrak{h}]$ -module is torsion-free, it follows that $I|_{\mathfrak{h}^{\text{reg}}} \neq 0$, where $I|_{\mathfrak{h}^{\text{reg}}}$ denotes the left (equivalently, right) localization of I viewed as a Harish-Chandra bimodule. We claim that $V|_{\mathfrak{h}^{\text{reg}}} = (\mathbf{H}_c|_{\mathfrak{h}^{\text{reg}}})/(I|_{\mathfrak{h}^{\text{reg}}}) = 0$. Indeed, as we have mentioned in §2, there is an algebra isomorphism $\mathbf{H}_c|_{\mathfrak{h}^{\text{reg}}} \simeq \mathcal{D}(\mathfrak{h}^{\text{reg}}) \# W$. The algebra $\mathcal{D}(\mathfrak{h}^{\text{reg}}) \# W$ is known to be simple [Mo]. Hence, $I|_{\mathfrak{h}^{\text{reg}}} \neq 0$ implies $\mathbf{H}_c|_{\mathfrak{h}^{\text{reg}}} = I|_{\mathfrak{h}^{\text{reg}}}$. Thus, $(\mathbf{H}_c/I)|_{\mathfrak{h}^{\text{reg}}} = 0$. Now, Corollary 3.6 yields: $0 = V = \mathbf{H}_c/I$, and $\mathbf{H}_c = I$. $\square$

From Proposition 3.5 one also obtains (see also [Gi]) the following important

Corollary 3.7. *If $c \in \mathbb{C}[R]_{\text{reg}}^W$, then every Harish-Chandra module has finite length.*

Proof. Let $V = V^0 \supsetneq V^1 \supsetneq \dots$ be a strictly decreasing chain of Harish-Chandra sub-bimodules of a Harish-Chandra bimodule V . View it as a chain of locally-free $\mathbb{C}[\mathfrak{h}]^W \otimes \mathbb{C}[\mathfrak{h}^*]^W$ -modules. Since each quotient V^i/V^{i+1} is a non-zero locally-free $\mathbb{C}[\mathfrak{h}]^W \otimes \mathbb{C}[\mathfrak{h}^*]^W$ -module (by Proposition 3.5), the ranks of the locally-free $\mathbb{C}[\mathfrak{h}]^W \otimes \mathbb{C}[\mathfrak{h}^*]^W$ -modules V^i form a strictly decreasing sequence: $\mathrm{rk}_{\mathbb{C}[\mathfrak{h}]^W \otimes \mathbb{C}[\mathfrak{h}^*]^W} V^0 > \mathrm{rk}_{\mathbb{C}[\mathfrak{h}]^W \otimes \mathbb{C}[\mathfrak{h}^*]^W} V^1 > \dots$. It follows that $V^N = 0$, for N big enough. $\square$

Let $\mathbf{x}^2 \in \mathbb{C}[\mathfrak{h}]$ denote the squared norm function: $x \mapsto (x, x)$, cf. §2, and let $\mathbf{y}^2 \in \mathbb{C}[\mathfrak{h}^*]$ denote a similar function on $\mathfrak{h}^*$ relative to the inner product transported from $\mathfrak{h}$ via the isomorphism $\mathfrak{h}^* \simeq \mathfrak{h}$. A straightforward computation based on formula (2.6) shows that the elements $\langle \mathbf{x}^2, \mathbf{h}, \mathbf{y}^2 \rangle$ form an $\mathfrak{sl}_2$ -triple in the algebra H_c .

Proposition 3.8. *Any Harish-Chandra bimodule V breaks, under the adjoint action of $\langle \mathbf{x}^2, \mathbf{h}, \mathbf{y}^2 \rangle$, into a direct sum of finite-dimensional $\mathfrak{sl}_2$ -modules. In particular, the $\mathrm{ad} \mathbf{h}$ -action on V is diagonalizable.*

Remark. We see that, in the situation of the Proposition the adjoint $\mathfrak{sl}_2$ -action exponentiates to an algebraic SL_2 -action on V . In particular, the automorphism corresponding to the action of the element $\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \in SL_2$ may be thought of as a sort of *Fourier transform*. In the special case $V = H_c$, this SL_2 -action on H_c by algebra automorphisms has been studied in [EG].

The SL_2 -action on H_c combined with SL_2 -action on a Harish-Chandra bimodule gives way to the following more symmetric point of view on Harish-Chandra bimodules.

Corollary 3.9. *A finitely generated $H_c \otimes (H_c)^{op}$ -module V is a Harish-Chandra bimodule if and only if, for any $g \in SL_2(\mathbb{C})$, the adjoint action on V of the subalgebra $g(\mathbb{C}[\mathfrak{h}]^W) \subset H_c$ is locally nilpotent.* $\square$

Write $U(-)$ for the enveloping algebra of a Lie algebra. We need the following lemma in representation theory of a complex semisimple Lie algebra $\mathfrak{g}$ with triangular decomposition: $\mathfrak{g} = \mathfrak{n}_+ + \mathfrak{h} + \mathfrak{n}_-$, which is a special case of a more general result due to Kac [K].

Lemma 3.10. *Let V be a $U(\mathfrak{g})$ -module such that, $\dim U(\mathfrak{n}_+)v < \infty$ and $\dim U(\mathfrak{n}_-)v < \infty$, for any $v \in V$. Then V is a locally finite $U(\mathfrak{g})$ -module.* $\square$

Proof of Proposition 3.8. We apply Lemma 3.10 to the Lie algebra $\mathfrak{g} = \mathfrak{sl}_2$ acting on a Harish-Chandra H_c -bimodule V via the adjoint action of the $\mathfrak{sl}_2$ -triple $\langle \mathbf{x}^2, \mathbf{h}, \mathbf{y}^2 \rangle$. The conditions of the Lemma clearly hold in this case. We conclude that the adjoint action on V of our $\mathfrak{sl}_2$ -triple is locally finite. The result follows. $\square$

4 The spherical subalgebra $eH_c e$

We begin with the following simple ring-theoretic result.

Lemma 4.1. *If A is a simple unital algebra and e is an idempotent of A . Then the algebra eAe is simple and Morita equivalent to A .*

Proof. Let I be a nonzero two-sided ideal in eAe . Then $AIA = A$ since A is simple, and hence $eAIAe = eAe$. But $I = eIe$, so $eAeIeAe = eAe$, which implies $I = eAe$. Hence eAe is simple. The second claim follows from the general fact that if $AeA = A$ then A is Morita equivalent to eAe (see e.g. [MR]). $\square$

Now, let W be our Coxeter group, and $\mathbf{e} = \frac{1}{|W|} \sum_{w \in W} w$ the symmetrizer in $\mathbb{C}W$. In [EG] we have introduced the subalgebra $\mathbf{e}H_c\mathbf{e} \subset H_c$, called the spherical subalgebra. From Lemma 4.1 and Theorem 3.1 we get

Corollary 4.2. *If $c \in \mathbb{C}[R]_{\text{reg}}^W$, then $\mathbf{e}H_c\mathbf{e}$ is a simple algebra, Morita equivalent to H_c .* $\square$

One defines the category $\mathcal{O}_{\mathbf{e}H_c\mathbf{e}}$ to be the abelian category formed by the finitely-generated $\mathcal{O}_{\mathbf{e}H_c\mathbf{e}}$ -modules M , such that the action on M of the subalgebra $\mathbb{C}[\mathfrak{h}^*]^W \subset \mathbf{e}H_c\mathbf{e}$ is locally finite.

From Corollary 4.2 we deduce

Proposition 4.3. *For any $c \in \mathbb{C}[R]_{\text{reg}}^W$, the functor $M \mapsto \mathbf{e} \cdot M$ gives an equivalence: $\mathcal{O}_{H_c} \xrightarrow{\sim} \mathcal{O}_{\mathbf{e}H_c\mathbf{e}}$.* $\square$

Put $M_{\mathbf{e}}(\lambda, \tau) := \mathbf{e}M(\lambda, \tau)$. This $\mathbf{e}H_c\mathbf{e}$ -module corresponds to the standard H_c -module via Morita equivalence. The modules $M_{\mathbf{e}}(\lambda, \tau)$ will be referred to as standard $\mathbf{e}H_c\mathbf{e}$ -modules. These modules can be explicitly described as follows. Given $\lambda \in \mathfrak{h}^*$, one has an algebra homomorphism: $\mathbb{C}[\mathfrak{h}^*]\#W_\lambda \twoheadrightarrow \mathbb{C}W_\lambda$ sending $P \in \mathbb{C}[\mathfrak{h}^*]$ to $P(\lambda)$ and identical on W_λ . We pull-back the left regular representation of W_λ to $\mathbb{C}[\mathfrak{h}^*]\#W_\lambda$ via this homomorphism, and form an induced H_c -module $N(\lambda) := \text{Ind}_{\mathbb{C}[\mathfrak{h}^*]\#W_\lambda}^{H_c} \mathbb{C}W_\lambda$. Observe further that the multiplication-action of the group W_λ on the group algebra $\mathbb{C}W_\lambda$ on the right commutes with the left $\mathbb{C}[\mathfrak{h}^*]\#W$ -module structure, hence, gives rise to a right W_λ -action on $N(\lambda) := \text{Ind}_{\mathbb{C}[\mathfrak{h}^*]\#W_\lambda}^{H_c} \mathbb{C}W_\lambda$. We leave to the reader to prove

Lemma 4.4. *The $\mathbf{e}H_c\mathbf{e}$ -module $M_{\mathbf{e}}(\lambda, \tau)$ is isomorphic to the τ -isotypic component (with respect to the right W_λ -action) in the left $\mathbf{e}H_c\mathbf{e}$ -module $\mathbf{e} \cdot N(\lambda)$.* $\square$

There is also a notion of Harish-Chandra bimodule over the spherical algebra $\mathbf{e}H_c\mathbf{e}$ instead of H_c , defined by the same Definition 3.2. An obvious analogue of Lemma 3.3 holds for Harish-Chandra bimodules over $\mathbf{e}H_c\mathbf{e}$. Further note that, for any $c \in \mathbb{C}[R]_{\text{reg}}^W$, all standard $\mathbf{e}H_c\mathbf{e}$ -modules $M_{\mathbf{e}}(\lambda, \tau)$ are simple, due to Morita equivalence of Proposition 4.3. Repeating the proof of Proposition 3.5 we deduce

Corollary 4.5. *If $c \in \mathbb{C}[R]_{\text{reg}}^W$, then any Harish-Chandra $\mathbf{e}H_c\mathbf{e}$ -bimodule V is a finite rank projective module over $\mathbb{C}[\mathfrak{h}]^W \otimes \mathbb{C}[\mathfrak{h}^*]^W \subset \mathbf{e}H_c\mathbf{e} \otimes (\mathbf{e}H_c\mathbf{e})^{op}$. In particular, $V|_{\mathfrak{h}^{\text{reg}}/W} = 0 \implies V = 0$.* $\square$

Remark. It has been explained in [EG, §8] that, in the case of a root system of type $\mathbf{A}_1$, the algebra $\mathbf{e}H_c\mathbf{e}$ is isomorphic to a quotient of the algebra $U(\mathfrak{sl}_2)$. Specifically, if $C = ef + fe + h^2/2 \in U(\mathfrak{sl}_2)$ is the standard quadratic Casimir, then $\mathbf{e}H_c\mathbf{e} \simeq U(\mathfrak{sl}_2)/I_c$, where I_c is the ideal in $U(\mathfrak{sl}_2)$ generated by the central element $C - \frac{1}{2}(c + \frac{3}{2}) \cdot (c - \frac{1}{2})$. Proposition 3.8 and Lemma 3.10 imply that a Harish-Chandra $\mathbf{e}H_c\mathbf{e}$ -bimodule is in this case nothing but a Harish-Chandra bimodule over $U(\mathfrak{sl}_2)$ (with equal³ left and right central characters), in the sense of representation theory of semisimple Lie algebras, see [Di]. Further, the isomorphism $\mathbf{e}H_c\mathbf{e} \simeq U(\mathfrak{sl}_2)/I_c$ suggests to view the standard module $M(\lambda, \sigma)$, $\lambda \neq 0$, as a generalized *Whittaker module*. $\square$

We now prove the following important result that reduces, in the special case $c = 0$ and $\mathbf{e}H_c\mathbf{e} = \mathcal{D}(\mathfrak{h})^W$, to Theorem 5 in the Levasseur-Stafford paper [LS] (for Coxeter groups).

Theorem 4.6 (Levasseur-Stafford theorem for $\mathbf{e}H_c\mathbf{e}$). *If the algebra $\mathbf{e}H_c\mathbf{e}$ is simple for some $c \in \mathbb{C}[R]^W$, then it is generated as an algebra by $\mathbb{C}[\mathfrak{h}]^W \cdot \mathbf{e}$ and $\mathbb{C}[\mathfrak{h}^*]^W \cdot \mathbf{e}$.*

³non-equal case will be treated in §8.

To prove the theorem we begin with the following *weak localized* version⁴ of a Poisson analogue of Theorem 5 of [LS].

Lemma 4.7. *The algebra $\mathbb{C}[\mathfrak{h}^{\text{reg}} \times \mathfrak{h}^*]^W$ of regular functions on the symplectic manifold $(\mathfrak{h}^{\text{reg}} \times \mathfrak{h}^*)/W$ is generated by $\mathbb{C}[\mathfrak{h}^{\text{reg}}]^W$ and $\mathbb{C}[\mathfrak{h}^*]^W$ as a Poisson algebra.*

Proof. (of Lemma 4.7) Denote by A the Poisson subalgebra of $\mathbb{C}[\mathfrak{h}^{\text{reg}} \times \mathfrak{h}^*]^W$, Poisson generated by $\mathbb{C}[\mathfrak{h}^{\text{reg}}]^W$ and $\mathbb{C}[\mathfrak{h}^*]^W$. By Hilbert-Noether lemma, A is a finitely generated commutative algebra without zero divisors. Let $Y = (\mathfrak{h}^{\text{reg}} \times \mathfrak{h}^*)/W$, and $Y' = \text{Spec}(A)$. Then Y' is an irreducible affine algebraic variety, and the natural map $f : Y \rightarrow Y'$ is a finite map. In particular, f is surjective.

Let us show that f is injective, i.e. that the algebra A separates points of Y . For this, consider the natural map $h : Y \rightarrow \mathfrak{h}^{\text{reg}}/W \times \mathfrak{h}^*/W$. Fix $(q, p) \in \mathfrak{h}^{\text{reg}} \times \mathfrak{h}^*$, let $W_p \subset W$ be the stabilizer of p , and $(\bar{q}, \bar{p})$ be the projection of (q, p) to $\mathfrak{h}^{\text{reg}}/W \times \mathfrak{h}^*/W$. The set $h^{-1}(\bar{q}, \bar{p})$ consists of equivalence classes of points (q, gp) , where $g \in W/W_p$. Thus, to prove the injectivity of f , it suffices to find an element $a \in A$ such that the numbers $a(q, gp)$ are distinct for all $g \in W/W_p$.

Let $z \in \mathfrak{h}^*$ be a vector such that the numbers $\langle z, gp \rangle$ are distinct for all $g \in W/W_p$ (this is the case for a generic vector, so such z exists). Let $b \in \mathbb{C}[\mathfrak{h}^{\text{reg}}]^W$ be a function such that $(db)(q) = z$ (such exists since the stabilizer of q is trivial). Set $a = \{b, \mathbf{y}^2/2\}$, where $\mathbf{y}^2$ is the squared norm function on $\mathfrak{h}^*$. Then $a(q, gp) = \langle z, gp \rangle$, which are distinct, as desired.

Thus, f is bijective. To conclude the proof, we need to show that moreover f is an isomorphism. Since $Y = (\mathfrak{h}^{\text{reg}} \times \mathfrak{h}^*)/W$ is smooth, it is enough to find, for any $(q_0, p_0) \in Y$, a set of elements of A which form a system of local coordinates near (q_0, p_0) .

Let $a_1, \dots, a_r \in \mathbb{C}[\mathfrak{h}^{\text{reg}}]^W$ be functions such that the vectors $da_i(q)$ form a basis of $\mathfrak{h}^*$. Let $b_i = \{a_i, \mathbf{y}^2/2\}$. It is easy to see by computing the Jacobian that the functions a_i and b_i form a system of coordinates on Y near (q_0, p_0) . The lemma is proved. $\square$

Lemma 4.8 ([LS], Lemma 9). *Let $R \subset S$ be two (not necessarily commutative) Noetherian domains, such that S is simple, and finite both as a left and a right R -module. Then, if R and S have the same skew field of fractions, then $R = S$.* $\square$

Proof of Theorem 4.6. The proof is analogous to the proof of Theorem 5 in [LS]. We apply Lemma 4.8 to the situation when $S = \mathbf{e}\mathcal{H}_c\mathbf{e}$, and R is the subalgebra generated by $\mathbb{C}[\mathfrak{h}]^{W \cdot \mathbf{e}}$ and $\mathbb{C}[\mathfrak{h}^*]^{W \cdot \mathbf{e}}$. It remains to check the conditions of the Lemma.

We know that $\mathbf{gr}(S) = \mathbb{C}[x, y]^W$ is a finite rank $\mathbb{C}[\mathfrak{h}^*]^W \otimes \mathbb{C}[\mathfrak{h}]^W$ module. Hence $\mathbf{gr}(S)$, and its submodule $\mathbf{gr}(R)$, are both Noetherian. Thus, so are R and S . It is also clear that R and S are domains (have no zero divisors). By assumption S is a simple algebra. Further, as we have seen above, $\mathbf{gr}(S)$ is a finite $\mathbf{gr}(R)$ module. Hence, S is a finite left and right R -module.

Finally, let us check that the quotient fields of R and S coincide. It suffices to check that $R|_{\mathfrak{h}^{\text{reg}}/W} = S|_{\mathfrak{h}^{\text{reg}}/W}$. Consider the filtration on $S|_{\mathfrak{h}^{\text{reg}}/W}$ defined by “degree in y ” (i.e. order of differential operators in the Dunkl-Cherednik realization). This filtration is nonnegative, so to check that $R|_{\mathfrak{h}^{\text{reg}}/W} = S|_{\mathfrak{h}^{\text{reg}}/W}$, it suffices to check that $\mathbf{gr}(R|_{\mathfrak{h}^{\text{reg}}/W}) = \mathbf{gr}(S|_{\mathfrak{h}^{\text{reg}}/W})$. The algebras $\mathbf{gr}(R|_{\mathfrak{h}^{\text{reg}}/W})$ and $\mathbf{gr}(S|_{\mathfrak{h}^{\text{reg}}/W})$ are Poisson algebras. Moreover, it is easy to see that $\mathbf{gr}(S|_{\mathfrak{h}^{\text{reg}}/W}) = \mathbb{C}[\mathfrak{h}^{\text{reg}} \times \mathfrak{h}^*]^W$, while $\mathbf{gr}(R|_{\mathfrak{h}^{\text{reg}}/W})$ contains $\mathbb{C}[\mathfrak{h}^{\text{reg}}]^W$ and $\mathbb{C}[\mathfrak{h}^*]^W$. Hence, it follows from Lemma 4.7 that $R|_{\mathfrak{h}^{\text{reg}}/W} = S|_{\mathfrak{h}^{\text{reg}}/W}$. Theorem 4.6 is proved. $\square$

We observe next that the elements $\langle \mathbf{x}^2\mathbf{e}, \mathbf{h}\mathbf{e}, \mathbf{y}^2\mathbf{e} \rangle$ form an $\mathfrak{sl}_2$ -triple in the algebra $\mathbf{e}\mathcal{H}_c\mathbf{e}$, and an analogue of Proposition 3.8 holds for Harish-Chandra $\mathbf{e}\mathcal{H}_c\mathbf{e}$ -bimodules. In particular, the adjoint

⁴the ‘strong’ version of the Lemma, i.e., the non-localized one where $\mathfrak{h}^{\text{reg}}$ is replaced by $\mathfrak{h}$, is known, due to Wallach [Wa], to be *false*, in general.

action of $\mathfrak{h}$ on a Harish-Chandra H_c -bimodule, resp., the adjoint $\mathfrak{h}\mathbf{e}$ -action on a Harish-Chandra $\mathbf{e}H_c\mathbf{e}$ -bimodule, V gives a grading $V = \bigoplus_{k \in \mathbb{Z}} V(k)$, where $V(k) = \{v \in V \mid \text{ad } \mathfrak{h}(v) = k \cdot v\}$. Therefore, the structure theory of finite dimensional representations of $\mathfrak{sl}_2$ implies that, for any $k \geq 0$ the operator $(\text{ad } \mathbf{y}^2)^k$ induces an isomorphism $(\text{ad } \mathbf{y}^2)^k : \text{Ker}(\text{ad } \mathbf{x}^2) \cap V(-k) \xrightarrow{\sim} \text{Ker}(\text{ad } \mathbf{y}^2) \cap V(k)$. In particular, for $V = H_c$, resp. $V = \mathbf{e}H_c\mathbf{e}$, the map $(\text{ad } \mathbf{y}^2)^k$ takes the space $\mathbb{C}^k[\mathfrak{h}] \subset H_c$ into the space formed by elements of filtration degree $\leq k$ and $\text{ad } \mathfrak{h}$ -weight $(-k)$, which is exactly the space $\mathbb{C}^k[\mathfrak{h}^*] \subset H_c$. Thus, we deduce

Corollary 4.9. *For any $k \geq 0$ the operator $(\text{ad } \mathbf{y}^2)^k$ induces isomorphisms*

$$\mathbb{C}^k[\mathfrak{h}] \xrightarrow{\sim} \mathbb{C}^k[\mathfrak{h}^*] \quad \text{and} \quad \mathbb{C}^k[\mathfrak{h}]^W \cdot \mathbf{e} \xrightarrow{\sim} \mathbb{C}^k[\mathfrak{h}^*]^W \cdot \mathbf{e} \quad , \quad \mathbb{C}[\mathfrak{h}], \mathbb{C}[\mathfrak{h}^*] \subset H_c. \quad \square$$

The natural $\mathbb{C}^*$ -action on $\mathfrak{h}^{\text{reg}}$ by dilations gives a $\mathbb{Z}$ -grading on $\mathcal{D}(\mathfrak{h}^{\text{reg}})$ by the *weight* of the $\mathbb{C}^*$ -action. Also, there is a standard filtration on $\mathcal{D}(\mathfrak{h}^{\text{reg}})$ by the *order* of differential operators, and we let $\mathcal{D}(\mathfrak{h}^{\text{reg}})_-$ denote the subset in $\mathcal{D}(\mathfrak{h}^{\text{reg}})$ spanned by all homogenous differential operators $u \in \mathcal{D}(\mathfrak{h}^{\text{reg}})$, such that $\text{weight}(u) + \text{order}(u) \leq 0$. Clearly, $\mathcal{D}(\mathfrak{h}^{\text{reg}})_-$ is an associative subalgebra in $\mathcal{D}(\mathfrak{h}^{\text{reg}})$.

Next, introduce the rational Calogero-Moser operator $L_c := \Delta - \sum_{\alpha \in R_+} \frac{2c_\alpha}{\alpha} \partial_\alpha$, where Δ denotes the 2-d order Laplacian on $\mathfrak{h}$. It is clear that $L_c \in \mathcal{D}(\mathfrak{h}^{\text{reg}})_-$. Following [O], we let $\mathcal{C}_c$ be the *centralizer* of L_c in the algebra $\mathcal{D}(\mathfrak{h}^{\text{reg}})_-^W = \mathcal{D}(\mathfrak{h}^{\text{reg}})^W \cap \mathcal{D}(\mathfrak{h}^{\text{reg}})_-$, and let $\mathcal{B}_c$ denote the subalgebra in $\mathcal{D}(\mathfrak{h}^{\text{reg}})^W$ generated by $\mathcal{C}_c$ and by $\mathbb{C}[\mathfrak{h}]^W$, the subalgebra of W -invariant polynomials.

Recall that in [EG] we have used the Dunkl representation of the algebra H_c to construct an injective algebra homomorphism $\Theta^{\text{spher}} : \mathbf{e}H_c\mathbf{e} \hookrightarrow \mathcal{D}(\mathfrak{h}^{\text{reg}})^W$ that sends the element $\mathbf{y}^2\mathbf{e}$ to L_c . The following result is a strengthening of [EG], Theorem 4.8.

Proposition 4.10. *For any $c \in \mathbb{C}[R]_{\text{reg}}^W$, the image of the map: $\mathbf{e}H_c\mathbf{e} \hookrightarrow \mathcal{D}(\mathfrak{h}^{\text{reg}})^W$ equals $\mathcal{B}_c$. Thus, one has an algebra isomorphism $\Theta^{\text{spher}} : \mathbf{e}H_c\mathbf{e} \xrightarrow{\sim} \mathcal{B}_c$.*

Proof. It is obvious that the map Θ^{spher} takes the subalgebra $\mathbb{C}[\mathfrak{h}]^W \cdot \mathbf{e} \subset \mathbf{e}H_c\mathbf{e}$ into $\mathbb{C}[\mathfrak{h}]^W \subset \mathcal{D}(\mathfrak{h}^{\text{reg}})^W$, and takes the subalgebra $\mathbb{C}[\mathfrak{h}^*]^W \cdot \mathbf{e} \subset \mathbf{e}H_c\mathbf{e}$ into $\mathcal{C}_c \subset \mathcal{D}(\mathfrak{h}^{\text{reg}})^W$. Hence, $\Theta^{\text{spher}}(\mathbf{e}H_c\mathbf{e}) \supset \mathcal{B}_c$. But, for $c \in \mathbb{C}[R]_{\text{reg}}^W$, the algebra $\mathbf{e}H_c\mathbf{e}$ is generated by $\mathbb{C}[\mathfrak{h}]^W \cdot \mathbf{e}$ and $\mathbb{C}[\mathfrak{h}^*]^W \cdot \mathbf{e}$, by the Levasseur-Stafford theorem. Thus, the algebra $\Theta^{\text{spher}}(\mathbf{e}H_c\mathbf{e})$ is generated by $\mathbb{C}[\mathfrak{h}]^W$ and $\mathcal{C}_c$. $\square$

Let $\varepsilon : W \rightarrow \mathbb{Z}/2\mathbb{Z}$ be a multiplicative character of the group W (if W is the Weyl group of a simply laced root system, then there is only one such character $\varepsilon(w) = \det_{\mathfrak{h}}(w)$). We consider the set: $R_\varepsilon := \{\alpha \in R \mid \varepsilon(s_\alpha) = -1\}$. Denote by $\mathbf{1}_\varepsilon$ the characteristic function of the subset $R_\varepsilon \subset R$, that is $\mathbf{1}_\varepsilon(\alpha) := 1$ if $\alpha \in R_\varepsilon$, and 0 otherwise. Clearly, $\mathbf{1}_\varepsilon \in \mathbb{C}[R]^W$. Also, we put $\mathbf{e}_\varepsilon := \frac{1}{|W|} \sum_{w \in W} \varepsilon(w) \cdot w$, a central idempotent in $\mathbb{C}W$.

Proposition 4.11. *There is an algebra isomorphism: $\mathbf{e}_\varepsilon H_c \mathbf{e}_\varepsilon \simeq \mathbf{e}H_{c-\mathbf{1}_\varepsilon} \mathbf{e}$, for any $c \in \mathbb{C}[R]_{\text{reg}}^W$.*

Proof. Since $c \in \mathbb{C}[R]_{\text{reg}}^W$, Corollary 4.9 implies that the algebra $\mathbf{e}H_c\mathbf{e}$ is generated by $\mathbb{C}[\mathfrak{h}]^W \cdot \mathbf{e}$ and by the element $\mathbf{y}^2\mathbf{e}$. Hence, the algebra $\Theta^{\text{spher}}(\mathbf{e}H_c\mathbf{e})$ is generated by $\mathbb{C}[\mathfrak{h}]^W$ and by the Calogero-Moser operator $\Theta^{\text{spher}}(\mathbf{y}^2\mathbf{e}) = L_c$. But $\Theta^{\text{spher}}(\mathbf{e}H_c\mathbf{e}) = \mathcal{B}_c$, by Proposition 4.10. Hence, $\mathcal{B}_c$ is generated by $\mathbb{C}[\mathfrak{h}]^W$ and L_c as an algebra.

Now, given a character $\varepsilon : W \rightarrow \{\pm 1\}$, we define a polynomial

$$\delta_\varepsilon = \prod_{\alpha \in R_+} \alpha^{\mathbf{1}_\varepsilon(\alpha)} = \prod_{\alpha \in R_\varepsilon \cap R_+} \alpha \in \mathbb{C}[\mathfrak{h}].$$

To emphasize dependence on the parameter ‘ c ’, write $\mathbf{y}^2 = \mathbf{y}_c^2 \in \mathbf{H}_c$, and $\Theta_c : \mathbf{H}_c \rightarrow \mathcal{D}(\mathfrak{h}^{\text{reg}}) \# W$, for the Dunkl representation. Then, an easy calculation, see [He], yields the following equation:

$$\delta_\varepsilon^{-1} \circ \Theta_c(\mathbf{y}_c^2 \mathbf{e}_\varepsilon) \circ \delta_\varepsilon = \Theta_{c-1_\varepsilon}(\mathbf{y}_{c-1_\varepsilon}^2 \mathbf{e}) \quad \text{holds in } \mathcal{D}(\mathfrak{h}^{\text{reg}}) \# W.$$

Since $e^{2\pi i c} = e^{2\pi i(c-1_\varepsilon)} \in \text{Reg}$, the algebra $\mathcal{B}_{c-1_\varepsilon} \simeq \mathbf{e} \mathbf{H}_{c-1_\varepsilon} \mathbf{e}$ is simple. Hence, it is generated by $\mathbb{C}[\mathfrak{h}]^W$ and $\mathbf{L}_{c-1_\varepsilon}$, and we obtain algebra isomorphisms:

$$\mathbf{e} \mathbf{H}_{c-1_\varepsilon} \mathbf{e} \simeq \mathcal{B}_{c-1_\varepsilon} = \delta_\varepsilon^{-1} \circ \Theta_c(\mathbf{e}_\varepsilon \mathbf{H}_c \mathbf{e}_\varepsilon) \circ \delta_\varepsilon \simeq \Theta_c(\mathbf{e}_\varepsilon \mathbf{H}_c \mathbf{e}_\varepsilon) \simeq \mathbf{e}_\varepsilon \mathbf{H}_c \mathbf{e}_\varepsilon. \quad \square$$

5 A trace on the Cherednik algebra

In this section we compute a trace on Cherednik algebras of type $\mathbf{A}_{n-1}$, that is on the algebra $\mathbf{H}_c$ corresponding to $W = S_n$, the Symmetric group. Write $[\mathbf{H}_c, \mathbf{H}_c]$ for the $\mathbb{C}$ -vector subspace in $\mathbf{H}_c$ spanned by all the commutators $[a, b]$, $a, b \in \mathbf{H}_c$. Let $\text{Tr}(g)$ denote the image of $g \in \mathbb{C}[S_n]$ under the composite map $\text{Tr} : \mathbb{C}[S_n] \hookrightarrow \mathbf{H}_c \twoheadrightarrow \mathbf{H}_c/[\mathbf{H}_c, \mathbf{H}_c]$. Given a permutation $g \in S_n$, write $\text{cycl}(g)$ for the number of cycles in g .

Proposition 5.1. *If $c \neq 0$ then, for any permutation $g \in S_n$, in $\mathbf{H}_c/[\mathbf{H}_c, \mathbf{H}_c]$ we have:*

$$\text{Tr}(g) = (1/nc)^{n-\text{cycl}(g)} \cdot \text{Tr}(1) \quad .$$

Proof. The proof proceeds by induction in $m = n - \text{cycl}(g)$. For $m = 0$, the result is obvious. Suppose the result is known for $m = p$, and let us establish it for $p + 1$. Let g have $n - p - 1$ cycles. Choose two indices $1 \leq i, j \leq n$, such that $g(i) = j$. Consider the permutation $\sigma = g \cdot s_{ij}$. Then $\sigma(j) = j$, and σ has $n - p$ cycles, so for σ the statement is known.

Let x_l, y_l be the elements in the first and second copy of $\mathfrak{h} \cong \mathfrak{h}^* \cong \mathbb{C}^{n-1} \subset \mathbb{C}^n$, respectively, corresponding to the vector $(-\frac{1}{n}, \dots, 1 - \frac{1}{n}, \dots, -\frac{1}{n})$, where $1 - \frac{1}{n}$ stands on the l -th place. The commutation relations in $\mathbf{H}_c$, see (1.1), imply readily that, for any l different from m , one has $[y_m, x_l] = c \cdot s_{ml} - \frac{1}{n}$. Thus, we find $[\sigma \cdot y_i, x_j] = \sigma \cdot [y_i, x_j] = c \cdot g - \frac{1}{n} \sigma$. Hence, $\text{Tr}(g) = \frac{1}{nc} \cdot \text{Tr}(\sigma)$, and we are done. $\square$

Recall that the (isomorphism classes of) irreducible representations of the Symmetric group are naturally labelled by Young diagrams. Write $Y(\tau)$ for the Young diagram (with n boxes) corresponding to a simple S_n -representation τ , and let s_τ denote the Schur function associated to $Y(\tau)$. Given a box $u \in Y(\tau)$ let $\text{cont}(u)$ denote the signed distance of u from the diagonal of $Y(\tau)$. We introduce a polynomial $F_\tau \in \mathbb{Z}[z]$ by the following two equivalent formulas

$$F_\tau(z) := \prod_{u \in Y(\tau)} (1 + \text{cont}(u)z) = h(\tau) \cdot s_\tau(1, z, \dots, z^{n-1}), \quad (5.2)$$

where $h(\tau)$ is the product of hook-lengths of elements of $Y(\tau)$ and $s_\tau(p_1, p_2, \dots, p_n)$ is the Schur polynomial of τ written in terms of power sums of symmetric variables. The equality of the two expressions in (5.2) follows from [Ma, Example 4, Ch. I].

Remark. The polynomial $F_\tau(z)$ has an interesting interpretation in terms of the Calogero-Moser space $\mathcal{M}_n$, cf. [Wi]. It is known that fixed points of a natural $\mathbb{C}^*$ -action on $\mathcal{M}_n$ are labelled by Young diagrams, cf. e.g. [EG, Prop. 4.16], [Ku]. Let (X_τ, Z_τ) denote a pair of $n \times n$ -matrices in the conjugacy class corresponding to the fixed point in $\mathcal{M}_n$ labelled by the Young diagram $Y(\tau)$ (for a direct construction of this point in terms of the S_n -module τ see [Go], and [EG, Conjecture 17.14]). Now, using formula [Wi, (6.14)] for Schur functions (and formula (2.3) in Appendix to [Wi]), one finds

$$F_\tau(z) = \det(-zX_\tau + (1 - Z_\tau)^{-1}).$$

Let $\mathbf{e}_\tau \in \mathbb{C}S_n \subset H_c$ denote the central idempotent in the group algebra corresponding to a simple S_n -module τ .

Theorem 5.3. *For any $c \in \mathbb{C}^*$ we have: $\text{Tr}(\mathbf{e}_\tau) = \frac{(\dim \tau)^2}{n!} \cdot F_\tau(\frac{1}{nc}) \cdot \text{Tr}(1)$.*

Proof. Let $\chi_\tau(g)$ denote the trace of $g \in S_n$ in the representation τ . Then one has an identity

$$\mathbf{e}_\tau = \frac{\dim \tau}{n!} \cdot \sum_{g \in S_n} \chi_\tau(g^{-1}) \cdot g. \quad (5.4)$$

To prove this, observe that both sides are central elements of $\mathbb{C}[S_n]$, and computing their traces in an arbitrary module $\sigma \in \text{Irrep}(S_n)$, we get

$$\frac{\dim \tau}{n!} \cdot \sum_{g \in S_n} \chi_\tau(g^{-1}) \cdot \chi_\sigma(g) = 1 \text{ if } \tau = \sigma \text{ and } 0 \text{ otherwise.}$$

The latter equation is nothing but the character orthogonality, and (5.4) follows. Therefore, using Proposition 5.1 we find

$$\text{Tr}(\mathbf{e}_\tau) = \frac{\dim \tau}{n!} \cdot \sum_{g \in S_n} \chi_\tau(g^{-1}) \cdot (1/nc)^{n-\text{cycl}(g)} \cdot \text{Tr}(1). \quad (5.5)$$

Now, using [S2, p.515, Exercise 7.50], we deduce: $\sum_{g \in S_n} \chi_\tau(g^{-1}) \cdot z^{n-\text{cycl}(g)} = \dim \tau \cdot F_\tau(z)$. $\square$

Remark. If the parameter $c \in \mathbb{C}^*$ is generic, then it has been shown in [EG] that the vector space $H_c/[H_c, H_c]$ is 1-dimensional. Moreover, we claim that for generic ‘ c ’ we have $\text{Tr}(1) \neq 0$. Indeed, assume the contrary, that is: $1 \in [H_c, H_c]$. This implies that there exists a positive integer $l = l(c)$ such that $1 \in [F_l H_c, F_l H_c]$, where $F_\bullet H_c$ is the standard increasing filtration on H_c . Note that for any given l , the set of ‘ c ’ such that $1 \in [F_l H_c, F_l H_c]$ is semialgebraic. Hence, one can find such an l that $1 \in [F_l H_c, F_l H_c]$, for all $c \in \mathbb{C}^*$ except possibly a finite set. But we will see below that there are infinitely many values of ‘ c ’ of the form $c = 1/n + \text{positive integer}$ such that the algebra H_c has a non-zero finite dimensional representation V . In that case the element $1 \in H_c$ acts as Id_V , hence has a nonzero trace. Therefore, one cannot have $1 \in [H_c, H_c]$, and our claim is proved.

Thus, choosing, for generic ‘ c ’, the element $\text{Tr}(1)$ as a basis in the 1-dimensional vector space $H_c/[H_c, H_c]$, one may view the map Tr as a ‘trace’: $H_c \rightarrow \mathbb{C}$.

We are going to apply the results above to study finite dimensional representations of H_c . Proposition 5.1 clearly yields

Corollary 5.6. *For any $g \in S_n$ and any finite dimensional H_c -module V , we have: $\text{Tr}|_V(g) = (1/nc)^{n-\text{cycl}(g)} \cdot \dim V$.* $\square$

Remark. For $c = \infty$, the Corollary implies that $V|_{S_n}$ is a multiple of the regular representation of S_n , which has been already shown in [EG].

Let $\mathbb{S}$ denote the set of all $c \in \mathbb{C}$ such that the algebra H_c (of type $\mathbf{A}_{n-1}$) has a nonzero finite dimensional representation. It is obvious from the above that $\mathbb{S}$ is a subset of $\mathbb{Q}^*$.

Proposition 5.7. *There is a subset $S \subset \{\frac{1}{n}, \frac{2}{n}, \dots, \frac{n-1}{n}\}$, such that $\mathbb{S} = -(S + \mathbb{Z}_{\geq 0}) \cup (S + \mathbb{Z}_{\geq 0})$.*

Proof. Throughout the proof we will freely exploit an isomorphism $H_c \simeq H_{-c}$ given by $s \mapsto -s$ for each simple reflection s , and $x \mapsto x, y \mapsto y$ for all $x \in \mathfrak{h}^*, y \in \mathfrak{h}$. In particular, we may assume $c > 0$, whenever necessary.

CLAIM 1. *If $c > 0$ is in $\mathbb{S}$ then either $c = p/n$ for some $p = 1, \dots, n-1$, or else $c > 1 - \frac{1}{n}$.*

To prove this, fix $c > 0$, and a non-zero finite dimensional H_c -module V . Using Theorem 5.3 we obtain the following formula for the multiplicity of occurrence of a simple S_n -module τ in V

$$[\tau : V] = \frac{1}{n!} \sum \chi_\tau(g^{-1}) \text{Tr}|_V(g) = \frac{\dim V \cdot \dim \tau}{n!} \cdot \prod_{u \in Y(\tau)} \left(1 + \frac{\text{cont}(u)}{nc}\right). \quad (5.8)$$

This means that the product $\prod_{u \in Y(\tau)} (1 + \text{cont}(u)/nc)$ has to be nonnegative for all τ . We deduce that either $c = p/n$, where $1 \leq p \leq n-1$ is a nonzero integer, or else $c > 1 - \frac{1}{n}$.

CLAIM 2. *Let $c \in \mathbb{S}$, and $c > 1 - \frac{1}{n}$. Then $c-1, c+1 \in \mathbb{S}$.*

To prove this, we first deduce from (5.8) that $[\text{triv} : V] \neq 0$. Thus eV is a nonzero $eH_c e$ -module.

Next, let $\varepsilon : S_n \rightarrow \{\pm 1\}$ be the sign character, and $e_\varepsilon \in \mathbb{C}[S_n]$ the corresponding central idempotent. Using that $c \geq 0$, we deduce similarly from (5.8) that $[\varepsilon : V] \neq 0$. It follows that $e_\varepsilon V$ is a nonzero $e_\varepsilon H_c e_\varepsilon$ -module.

Now, Proposition 4.11 implies that the algebras $eH_c e$ and $e_\varepsilon H_c e_\varepsilon$ are isomorphic to $e_\varepsilon H_{c+1} e_\varepsilon$ and $eH_{c-1} e$, respectively. Thus, the latter algebras have non-zero finite dimensional representations. Hence, so do H_{c+1}, H_{c-1} (by taking induced module from the corresponding spherical subalgebra). Claim 2 is proved.

By a similar argument one proves

- *Let $0 < c \leq 1 - \frac{1}{n}$. If H_c has a non-zero finite dimensional representation then so does H_{c+1} .*

Now we complete the proof of the Proposition. Fix an element $c \in \mathbb{S}$ such that $c > 0$. If $c > 1 - 1/n$ then $c \geq 1 + 1/n$, since in this case by Claim 2, $c-1$ should be in $\mathbb{S}$, while we know from Claim 1 that $\mathbb{S}$ has no intersection with the open interval $(-1/n, 1/n)$. So let c' be the fractional part of c . Then c' is in $\mathbb{S}$, by Claim 2. Let $S = \mathbb{S} \cap \{1/n, 2/n, \dots, 1 - 1/n\}$. Then $c' \in S$, so $\mathbb{S} \subset -(S + \mathbb{Z}_{\geq 0}) \cup (S + \mathbb{Z}_{\geq 0})$. The fact that this inclusion is an equality follows from the statement above that for $c > 0$, if $c \in \mathbb{S}$ then $c+1 \in \mathbb{S}$. $\square$

Remark. Note that $1/n \in \mathbb{S}$ since for $c = 1/n$ there exists an action of H_c on the trivial representation of S_n , with x_i and y_i acting by zero.

Proposition 5.1 yields a nice closed expression for the character of S_n -action on any finite dimensional H_c -module, up to a constant factor. To explain this, write $\mathfrak{h}_z := \mathbb{Z}^{n-1} \subset \mathbb{Z}^n$ for the set of n -tuples of integers with total sum zero, a $\mathbb{Z}$ -lattice in $\mathfrak{h}$. The group S_n acts naturally on $\mathfrak{h}_z$ by permutation. This induces, for any integer $r > 0$, an S_n -action on the finite set $\mathfrak{h}_z/r \cdot \mathfrak{h}_z \subset \mathbb{Z}^n/r \cdot \mathbb{Z}^n$, making the $\mathbb{C}$ -vector space $\mathbb{C}[\mathfrak{h}_z/r \cdot \mathfrak{h}_z]$ an S_n -module.

Let $K(S_n)$ be the Grothendieck group of finite dimensional representation of S_n . Given a finite dimensional H_c -module V , let $[V|_{S_n}] \in K(S_n)$ denote the class of its restriction to S_n . Note further that $V \neq 0$ implies, by Proposition 5.7, that $cn = r$ is an integer, not divisible by n .

Theorem 5.9. *Assume $c = r/n > 0$, and let V be a finite dimensional H_c -module.*

- (i) *There exists an integer $\ell > 0$ such that in $K(S_n)$ we have: $[V|_{S_n}] = \frac{\ell}{n} \cdot \text{class of } \mathbb{C}[(\mathbb{Z}/r\mathbb{Z})^n]$.*
- (ii) *Moreover, if r is coprime to n then $[V|_{S_n}] = \ell \cdot \text{class of } \mathbb{C}[\mathfrak{h}_z/r \cdot \mathfrak{h}_z]$.*

Proof. To prove (i) it suffices to show that the characters of S_n representations on both sides are proportional. But it is straightforward to see that the character of S_n -action on $\mathbb{C}[(\mathbb{Z}/r\mathbb{Z})^n]$ is given, by $\text{Tr}(g) = r^{\text{cyc1}(g)} = (cn)^{\text{cyc1}(g)}$, which up to a scalar is exactly the formula of Proposition 5.1. The integrality of the coefficient ℓ in part (i) follows from the fact that ℓ is equal to the trace of the cyclic permutation in $(\ell/n) \cdot \mathbb{C}[(\mathbb{Z}/r\mathbb{Z})^n]$. Part (ii) readily follows from (i) by comparing characters. $\square$

Conjecture 5.10. Let $c = r/n > 0$. The algebra H_c has a nonzero finite dimensional representation V if and only if r is coprime to n , moreover, in such a case in Theorem 5.9(ii) we have $\ell = 1$.

This conjecture has been proved in [BEG1].

6 The $eH_c e$ -module structure on quasi-invariants

In this section we review the definition and prove some basic properties of algebras of quasi-invariants of finite reflection groups. Most of these results have already appeared in the literature (see [CV], [CV2], [VSC], [FV], [EG2]).

Let $\mathcal{H}ol(U)$ be the space of holomorphic functions on an open set $U \subset \mathfrak{h}$, and $\mathcal{H}ol(\mathfrak{h})^W \subset \mathcal{H}ol(\mathfrak{h})$, the space of W -invariant holomorphic functions on $\mathfrak{h}$. Observe that $f \in \mathcal{H}ol(\mathfrak{h})$ is W -invariant if and only if $s_\alpha(f) = f$ for all $\alpha \in R_+$, since the set $\{s_\alpha \mid \alpha \in R_+\} \subset W$ generates W . We will extend the algebra $\mathcal{H}ol(\mathfrak{h})^W$ by relaxing the notion of reflection invariance in the following way. Let $m \in \mathbb{Z}_+$ be a non-negative integer, $s_\alpha \in W$ a reflection, and $x \in \mathfrak{h}$ a point on the corresponding hyperplane, i.e., $s_\alpha(x) = x$. We say that a holomorphic function f on an s_α -stable open neighborhood $U_x \ni x$ is *m-quasi-invariant* with respect to s_α if

$$s_\alpha(f) \equiv f \pmod{(\alpha)^{2m+1}}, \quad (6.1)$$

where (α) stands for the (principal) ideal in $\mathcal{H}ol(U_x)$ generated by the linear form α . To compare, a function $f \in \mathcal{H}ol(U_x)$ being invariant under s_α means that all *odd* powers of α in the Taylor expansion of f in the direction α vanish, while $f \in \mathcal{H}ol(U_x)$ being *m-quasi-invariant* means that such powers vanish only up to order $2m - 1$ (inclusive).

Let $\mathbb{Z}(R)_+^W$ be the set of all W -invariant functions on the root system R with values in non-negative integers.

Definition 6.2 (cf. [CV], [CV2]). Given $c \in \mathbb{Z}(R)_+^W$, a holomorphic function $f \in \mathcal{H}ol(\mathfrak{h})$ is said to be *c-quasi-invariant* with respect to W if, for each $\alpha \in R$, one has $s_\alpha(f) \equiv f \pmod{(\alpha)^{2c_\alpha+1}}$.

Given $c \in \mathbb{Z}(R)_+^W$, let Q_c denote the set of all *c-quasi-invariant polynomials* with respect to W . The following lemma collects some basic algebraic properties of quasi-invariants.

Lemma 6.3. (i) *The vector space Q_c is W -stable, and we have: $\mathbb{C}[\mathfrak{h}]^W \subseteq Q_c \subseteq \mathbb{C}[\mathfrak{h}]$.*

(ii) *Q_c is a finitely generated graded subalgebra of $\mathbb{C}[\mathfrak{h}]$, such that the integral closure of Q_c equals $\mathbb{C}[\mathfrak{h}]$.*

Proof. (i) is clear. Further, if $p, q \in \mathbb{C}[\mathfrak{h}]$ satisfy (6.1) with $m = c_\alpha$, for some $\alpha \in R_+$, so do their linear combination and product: for example, $s_\alpha(pq) = s_\alpha(p)s_\alpha(q) \in pq + (\alpha)^{2c_\alpha+1}$. Now, if (6.1) holds for all $\alpha \in R_+$ and $\beta \in R_+$ then $s_\alpha[s_\beta(p)] = s_\beta[s_{s_\beta(\alpha)}(p)] \in s_\beta(p) + (\alpha)^{2c_{s_\beta(\alpha)}+1} = s_\beta(p) + (\alpha)^{2c_\alpha+1}$, where the last equality holds due to the W -invariance of c . Thus Q_c is a subalgebra of $\mathbb{C}[\mathfrak{h}]$ invariant under the action W . Since W is a finite group, $\mathbb{C}[\mathfrak{h}]$ is finite as an $\mathbb{C}[\mathfrak{h}]^W$ -module, see e.g. [S], hence, as a Q_c -module. Then finiteness of Q_c as a $\mathbb{C}$ -algebra follows from the Hilbert-Noether Lemma (see [AM], Proposition 7.8). Further, $\mathbb{C}[\mathfrak{h}]$ being finite as a module over Q_c is equivalent to $\mathbb{C}[\mathfrak{h}]$ being integral as a ring over Q_c . On the other hand, the subalgebra Q_c contains an ideal of $\mathbb{C}[\mathfrak{h}]$, namely a sufficiently high power of $\delta = \prod_{\alpha \in R} \alpha$. Hence, Q_c and $\mathbb{C}[\mathfrak{h}]$ have the same field of fractions. It follows that $\mathbb{C}[\mathfrak{h}]$ is the integral closure of Q_c . $\square$

We recall the following elementary but basic observation [C],[FV]:

Lemma 6.4. *The action of the Calogero-Moser operator L_c preserves the subspace $Q_c \subset \mathbb{C}[\mathfrak{h}^{\text{reg}}]$, i.e., $L_c(Q_c) \subseteq Q_c$. $\square$*

Recall that according to Opdam [O], for any $c \in \mathbb{C}[R]^W$, there is an algebra isomorphism $\mathbb{C}[\mathfrak{h}^*]^W \xrightarrow{\sim} \mathcal{C}_c$, where $\mathcal{C}_c$ stands for the centralizer in $\mathcal{D}(\mathfrak{h}^{\text{reg}})^W$ of the Calogero-Moser operator L_c . Identify $\mathbb{C}[\mathfrak{h}^*]^W$ with $\mathbb{C}[\mathfrak{h}]^W$ via an invariant form on $\mathfrak{h}$. In [CV], Chalykh-Veselov extended Opdam's isomorphism: $\mathbb{C}[\mathfrak{h}]^W = \mathbb{C}[\mathfrak{h}^*]^W \xrightarrow{\sim} \mathcal{C}_c$ to an injective algebra homomorphism: $Q_c \hookrightarrow \mathcal{D}(\mathfrak{h}^{\text{reg}})^W$, $P \mapsto \mathcal{L}_P$. There is an explicit formula for the restriction of the map $P \mapsto \mathcal{L}_P$ to the subspace $Q_c \cap \mathbb{C}^k[\mathfrak{h}^*]$ of homogeneous polynomials in Q_c of degree $k \geq 1$. Specifically, according to [Be], cf. also §7 below, one has

$$\mathcal{L}_P = \frac{1}{2^k k!} (\text{ad } L_c)^k(P), \quad (6.5)$$

where P is a homogeneous polynomial in Q_c of degree $k \geq 1$. In the special case when $P \in \mathbb{C}^k[\mathfrak{h}^*]^W$ is W -invariant, this formula is an immediate (up to computing the constant factor, $\frac{1}{2^k k!}$, which is also easy) consequence of Corollary 4.9, which yields the isomorphism $\Theta^{\text{spher}}(\text{ad } \mathbf{y}^2)^k : \Theta^{\text{spher}}(\mathbb{C}^k[\mathfrak{h}]^W \cdot \mathbf{e}) \xrightarrow{\sim} \Theta^{\text{spher}}(\mathbb{C}^k[\mathfrak{h}^*]^W \cdot \mathbf{e})$.

Let $Q_c^b \subset \mathcal{D}(\mathfrak{h}^{\text{reg}})^W$ denote the image of the Chalykh-Veselov homomorphism $P \mapsto \mathcal{L}_P$. Thus Q_c^b is a commutative subalgebra in $\mathcal{D}(\mathfrak{h}^{\text{reg}})^W$ isomorphic to Q_c . As has been observed in [FV], formula (6.5) and Lemma 6.4 imply that the action on $\mathbb{C}[\mathfrak{h}^{\text{reg}}]^W$ of the algebra Q_c^b preserves the space Q_c , that is, $u(Q_c) \subset Q_c$, for any $u \in Q_c^b$. In particular, $\mathcal{C}_c(Q_c) \subset Q_c$. Thus, there is a natural action on Q_c of the algebra $\mathcal{B}_c \subset \mathcal{D}(\mathfrak{h}^{\text{reg}})^W$, generated by $\mathcal{C}_c$ and $\mathbb{C}[\mathfrak{h}]^W = Q_c^W$. This $\mathcal{B}_c$ -action clearly commutes with the W -action on Q_c . Thus, since $\mathcal{B}_c = \Theta^{\text{spher}}(\mathbf{eH}_c\mathbf{e})$ by Proposition 4.10, the space Q_c acquires an $\mathbb{C}W \otimes \mathbf{eH}_c\mathbf{e}$ -module structure. The structure of this module is completely described by the following proposition.

For any irreducible representation τ of W , let τ'_c be the representation of W for which the monodromy representation of the Dunkl connection with values in τ'_c is τ (see [O1] and references therein).⁵

Proposition 6.6. *There is a $\mathbb{C}W \otimes \mathbf{eH}_c\mathbf{e}$ -module isomorphism: $Q_c = \bigoplus_{\tau \in \text{Irrrep}(W)} \tau^* \otimes M_{\mathbf{e}}(\tau'_c)$.*

Proof. First of all, Q_c is finitely generated over $\mathbb{C}[\mathfrak{h}]^W$, hence over $\mathbf{eH}_c\mathbf{e}$. Further, the action of any homogeneous element of Q_c^b lowers the degree of polynomials in Q_c . Hence, the action on Q_c of the augmentation ideal of the algebra $\mathbb{C}[\mathfrak{h}^*]^W \subset \mathbf{eH}_c\mathbf{e}$ is locally nilpotent. It follows that $Q_c \in \mathcal{O}_{\mathbf{eH}_c\mathbf{e}}(0)$. Hence, Lemma 2.5(iv) implies that Q_c has finite length, as a $\mathbf{eH}_c\mathbf{e}$ -module. Recall further that, for any $c \in \mathbb{Z}[R]^W_+$, the standard modules $M(\tau)$ are irreducible. By Morita equivalence of H_c and $\mathbf{eH}_c\mathbf{e}$, for any τ , the modules $M_{\mathbf{e}}(\tau) := \mathbf{e}M(\tau)$ over $\mathbf{eH}_c\mathbf{e}$ are also irreducible. Thus, Q_c has a finite composition series whose simple subquotients are standard modules over $\mathbf{eH}_c\mathbf{e}$.

Observe next that the direct sum decomposition of the Proposition is easily checked directly for $c = 0$. Now, consider the localized algebra $(\mathbf{eH}_c\mathbf{e})|_{\mathfrak{h}_{\text{reg}}/W}$ and the module $Q_c|_{\mathfrak{h}_{\text{reg}}/W}$ over this algebra. It is clear from the existence of the Dunkl representation that both the localized algebra and module are independent of c , i.e. are the same as those for $c = 0$ (namely, they are the algebra of differential operators on $\mathfrak{h}_{\text{reg}}/W$ and the D-module on $\mathfrak{h}_{\text{reg}}/W$ corresponding to the

⁵A priori, the monodromy representation is a representation of the braid group of W , but since c is integer-valued, it factors through W . It is known ([O1]) that $\tau'_c = \tau$ for all irreducible Coxeter groups except E_7, E_8, H_3, H_4 , and even dihedral groups; otherwise $\tau'_c = \gamma^c(\tau)$, where γ is a certain involution on the set of irreducible representations of W , and for even dihedral groups γ^c stands for $\gamma^{c_1+c_2}$. Also, it is known that characters of τ and τ'_c coincide on reflections.

regular representation of W). Therefore, we have $Q_c|_{\mathfrak{h}^{\text{reg}}/W} = \bigoplus_{\tau \in \text{Irrep}(W)} \tau^* \otimes M_{\mathbf{e}}(\tau'_c)|_{\mathfrak{h}^{\text{reg}}/W}$ (by the definition of τ'_c).

Now, define $M'_{\mathbf{e}}(\tau) := (\tau \otimes Q_c)^W$. This is an $\mathbf{eH}_c\mathbf{e}$ -module, such that the localized module $M'_{\mathbf{e}}(\tau)|_{\mathfrak{h}^{\text{reg}}/W}$ coincides with $M_{\mathbf{e}}(\tau'_c)|_{\mathfrak{h}^{\text{reg}}/W}$. On the other hand, $M'_{\mathbf{e}}(\tau) \in \mathcal{O}_{\mathbf{eH}_c\mathbf{e}}(0)$, so it has a finite Jordan-Hölder series $0 = F_0 \subset F_1 \subset \dots \subset F_l = M'_{\mathbf{e}}(\tau)$, with simple subquotients F_i/F_{i-1} being isomorphic to standard modules $M_{\mathbf{e}}(\sigma_i)$, $i = 1, \dots, l$, see Proposition 4.3. The localization functor being exact, we see that $M'_{\mathbf{e}}(\tau)|_{\mathfrak{h}^{\text{reg}}/W}$ has a composition series with the subquotients $(F_i/F_{i-1})|_{\mathfrak{h}^{\text{reg}}/W} = M_{\mathbf{e}}(\sigma_i)|_{\mathfrak{h}^{\text{reg}}/W}$. But since $M'_{\mathbf{e}}(\tau)|_{\mathfrak{h}^{\text{reg}}/W} = M_{\mathbf{e}}(\tau'_c)|_{\mathfrak{h}^{\text{reg}}/W}$, the latter composition series must consist of a single term, i.e., $l = 1$. Thus, we find that $M'_{\mathbf{e}}(\tau) = M_{\mathbf{e}}(\tau'_c)$, and we are done. $\square$

Given a diagonalisable endomorphism A on a vector space E with finite-dimensional eigenspaces E_a , $a \in \mathbb{C}$, we put $\text{Tr}|_E(t^A) := \sum_{a \in \mathbb{C}} \dim_{\mathbb{C}} E_a \cdot t^a$, viewed as a formal infinite sum. For example, let eu denote the Euler operator on $\mathbb{C}[\mathfrak{h}]$ such that $\text{eu}(P) = (\deg P) \cdot P$, for any homogeneous polynomial P . Write $\chi_{\tau}(t) = \text{Tr}|_{(\tau \otimes \mathbb{C}[\mathfrak{h}])^W}(t^{\text{eu}})$ for the Poincaré series of the graded space $(\tau \otimes \mathbb{C}[\mathfrak{h}])^W$. These series are classically known.

Let $\kappa_c = \frac{1}{2} \sum_{\alpha \in R} c_{\alpha}(1 - s_{\alpha})$ be the central element of $\mathbb{C}W$ canonically attached to $c \in \mathbb{C}[R]^W$. This element acts by a scalar, say $\kappa_c(\tau)$, in each irreducible representation $\tau \in \text{Irrep}(W)$. We have

$$\text{Tr}|_{M_{\mathbf{e}}(\tau)}(t^{\mathbf{h}}) = t^{\frac{\dim \mathfrak{h}}{2} - \frac{1}{2} \sum_{\alpha \in R} c_{\alpha} + \kappa_c(\tau)} \cdot \chi_{\tau}(t). \quad (6.7)$$

To prove this formula, one first finds the eigenvalue of $\mathbf{h}$ -action on the "top" subspace $\tau \subset M(\tau)$. A straightforward calculation gives

$$\mathbf{h}|_{\tau} = \left(\frac{\dim \mathfrak{h}}{2} - \frac{1}{2} \sum_{\alpha \in R} c_{\alpha} + \kappa_c(\tau) \right) \cdot \text{Id}_{\tau}.$$

Further, it is easy to see from the Dunkl representation that $\mathbf{h}|_{\mathbb{C}[\mathfrak{h}]} = \text{eu} + \frac{\dim \mathfrak{h}}{2} - \frac{1}{2} \sum_{\alpha \in R} c_{\alpha}$. Formula (6.7) follows. $\square$

Applying Proposition 6.6, we obtain a formula for the Poincaré series of Q_c , i.e. for the function $\text{Tr}|_{Q_c}(t^{\text{eu}})$. The same formula has been found by Felder and Veselov ([FeV]), using a different method.

Corollary 6.8. $\text{Tr}|_{Q_c}(t^{\text{eu}}) = \sum_{\tau \in \text{Irrep}(W)} \dim \tau \cdot t^{\kappa_c(\tau)} \cdot \chi_{\tau}(t).$ $\square$

Proof. This follows from Proposition 6.6 and the equality that $\kappa_c(\tau) = \kappa_c(\tau'_c)$, which follows from the fact that the characters of τ and τ'_c coincide on reflections. $\square$

7 Differential operators on quasi-invariants

We recall some basic results and notation regarding differential operators on algebraic varieties. Let A be a commutative $\mathbb{C}$ -algebra, and let M and N be a pair of A -modules. The space $\mathcal{D}(M, N)$ of linear differential operators from M to N over A is a filtered subspace $\mathcal{D}(M, N) = \bigcup_{n \geq 0} \mathcal{D}_n(M, N) \subset \text{Hom}_{\mathbb{C}}(M, N)$ defined inductively by $\mathcal{D}_{-1}(M, N) = 0$, and

$$\mathcal{D}_n(M, N) := \{u \in \text{Hom}_{\mathbb{C}}(M, N) \mid u \circ a - a \circ u \in \mathcal{D}_{n-1}(M, N) \text{ for all } a \in A\}, \quad n > 0. \quad (7.1)$$

The elements of $\mathcal{D}_n(M, N) \setminus \mathcal{D}_{n-1}(M, N)$ are called *differential operators of order n* .

If $N = M = A$, we write $\mathcal{D}(A) := \mathcal{D}(A, A)$, which is clearly an algebra over $\mathbb{C}$. In particular, if $A = \mathbb{C}[X]$ is the ring of regular functions on an irreducible affine algebraic variety X , then $\mathcal{D}(A)$ is nothing but $\mathcal{D}(X)$, the algebra of differential operators on X .

If X is singular, the structure of $\mathcal{D}(X)$ can be quite complicated and is not well understood. A natural approach to study the algebra $\mathcal{D}(X)$ on an arbitrary (irreducible) variety X is to compare it with the ring of differential operators on the normalization of X (see [SS]). In more detail, let $\pi : \tilde{X} \rightarrow X$ be the normalization map. Write $\mathbb{K} := \mathbb{C}(X)$ for the field of rational functions on X (so that $\mathbb{C}[X] \subseteq \mathbb{C}[\tilde{X}] \subset \mathbb{K}$, and $\mathbb{C}[\tilde{X}]$ is the integral closure of $\mathbb{C}[X]$ in $\mathbb{K}$) and denote by $\mathcal{D}(\mathbb{K})$ the algebra of differential operators over $\mathbb{K}$. Since $\mathcal{D}(\mathbb{K}) \cong \mathbb{K} \otimes_{\mathbb{C}[X]} \mathcal{D}(X)$, we can identify $\mathcal{D}(X)$ with its image in $\mathcal{D}(\mathbb{K})$. This way we get

$$\mathcal{D}(X) = \{ u \in \mathcal{D}(\mathbb{K}) \mid u(\mathbb{C}[X]) \subseteq \mathbb{C}[X] \} \quad , \quad \mathcal{D}(\tilde{X}) = \{ u \in \mathcal{D}(\mathbb{K}) \mid u(\mathbb{C}[\tilde{X}]) \subseteq \mathbb{C}[\tilde{X}] \} . \quad (7.2)$$

Thus, both $\mathcal{D}(X)$ and $\mathcal{D}(\tilde{X})$ can be viewed as subalgebras in $\mathcal{D}(\mathbb{K})$.

From now on, we fix $c \in \mathbb{Z}[R]_+^W$ and consider the scheme $X_c := \text{Spec } Q_c$. Given $x \in \mathfrak{h}$, let $W_x \subset W$ denote the stabilizer of x under the natural action of W on $\mathfrak{h}$.

- Lemma 7.3.** (i) X_c is a rational irreducible affine algebraic variety with W -action. The normalization of X_c is isomorphic to $\mathfrak{h}$.
(ii) The normalization map $\pi_c : \mathfrak{h} \twoheadrightarrow X_c$ is bijective.
(iii) The local ring $\mathcal{O}_x(X_c)$ at a point $x \in X_c$ can be identified with the subring of $\mathbb{C}(\mathfrak{h})$ consisting of rational functions on $\mathfrak{h}$ regular at $\pi_c^{-1}(x)$ and quasi-invariant under W_x .

Proof. Part (i) is a reformulation of Lemma 6.3. To prove (ii) we only need to check injectivity of the normalization map. Thus, we have to show that for any points $x, x' \in \mathfrak{h}$, $x' \neq x$, there exists a quasi-invariant polynomial $p \in Q_c$ such that $p(x) \neq p(x')$. To this end, fix $x \in \mathfrak{h}$ and put

$$\delta_{2c+1,x} = \prod_{\{\alpha \in R_+ \mid (\alpha, x) \neq 0\}} \alpha^{2c_\alpha+1}, \quad \text{this is a quasi-invariant polynomial under all reflections in } W \setminus W_x, \text{ and also invariant under } W_x. \quad (7.4)$$

Now, given $x' \neq x$, choose any polynomial g such that $g(x) \neq 0$, and $g(x') = 0$. Define the polynomial $p(z) := \delta_{2c+1,x}(z) \cdot \prod_{w \in W_x} g(wz)$. We have: $p(x) \neq 0$, and $p(x') = 0$. Further, p is invariant under W_x and quasi-invariant under all reflections in $W \setminus W_x$, by construction. Thus, p is quasi-invariant and (ii) is proved.

To prove (iii) observe that $\mathcal{O}_x(X_c) \subset \{r \in \mathbb{C}(\mathfrak{h}) \mid r \text{ is regular at } x\}$. Conversely, let r be a rational function regular at x which is also quasi-invariant with respect to the group W_x . We can write $r = p/q$, where p, q are polynomials, and $q(x) \neq 0$. To show that $r = p/q \in \mathcal{O}_x(X_c)$ we set $f = \prod_{\{w \in W_x \mid w \neq 1\}} w(q)$, and write $r = \frac{p}{q} = \frac{pf\delta_{2c+1,x}}{qf\delta_{2c+1,x}}$, where $\delta_{2c+1,x}$ was defined in (7.4). The denominator in the last fraction is W_x -invariant, and also quasi-invariant under all the reflections which are not in W_x by construction. Thus, the denominator is quasi-invariant under W and nonvanishing at x . The numerator is also quasi-invariant, since so is the whole function r . Thus, $r \in \mathcal{O}_x(X_c)$, as desired. $\square$

According to (7.2) we may identify the algebra $\mathcal{D}(X_c)$ with the set $\{u \in \mathcal{D}(\mathfrak{h}^{\text{reg}}) \mid u(Q_c) \subset Q_c\}$. In the previous section we have introduced the subalgebra $\mathcal{D}(\mathfrak{h}^{\text{reg}})_- = \text{span of } \{u \in \mathcal{D}(\mathfrak{h}^{\text{reg}}) \mid \text{weight}(u) + \text{order}(u) \leq 0\}$. Put $\mathcal{D}(X_c)_- := \mathcal{D}(X_c) \cap \mathcal{D}(\mathfrak{h}^{\text{reg}})_-$. Recall further the subalgebra $Q_c^b \subset \mathcal{D}(\mathfrak{h}^{\text{reg}})_-$, the image of Q_c under the Chalykh-Veselov homomorphism $P \mapsto \mathcal{L}_P$.

Theorem 7.5. (i) We have: $\mathcal{D}(X_c)_- = Q_c^b$. This is a maximal commutative subalgebra in $\mathcal{D}(X_c)$, equal to the centralizer in $\mathcal{D}(\mathfrak{h}^{\text{reg}})_-$ of the Calogero-Moser operator L_c .

(ii) The left multiplication by Q_c and right multiplication by Q_c^b make $\mathcal{D}(X_c)$ into a rank one projective $Q_c \otimes Q_c^b$ -module.

The rest of the section is effectively devoted to the proof of this Theorem. The argument exploits crucially the notion of the Baker-Akhiezer (holomorphic) function $\psi_c : (x, k) \mapsto \psi_c(x, k) = \Phi(x, k) \cdot e^{(x, k)}$, where $x \in \mathfrak{h}$, $k \in \mathfrak{h}^*$, and Φ is a polynomial on $\mathfrak{h} \times \mathfrak{h}^*$, see [CV, FV, EG2].

Whenever necessary we will identify below $\mathfrak{h}^*$ with $\mathfrak{h}$ without further comment, using some fixed W -invariant form. Given $x \in \mathfrak{h}$, put

$$Q_{c,x} := \{f \in \mathbb{C}[\mathfrak{h}] \mid k \mapsto f(k)e^{kx} \text{ is a quasi-invariant holomorphic function}\}.$$

Proposition 7.6. For any $x \in \mathfrak{h}$, the space $Q_{c,x}$ is a projective rank one Q_c -module (not free, in general).

Proof. Fix $x_0 \in \mathfrak{h}$. We will show that the module Q_{c,x_0} is locally-free in a neighborhood of any point $k_0 \in \mathfrak{h}$. Let $g(t, k) := \psi_c(t \cdot x_0, k)$, where $t \in \mathbb{C}$ is a complex variable. Then, for each $k \in \mathfrak{h}$, the map: $t \mapsto g(t, k)$ is an entire function on $\mathbb{C}$. Write $g(t, k) = \sum_{n=0}^{\infty} g_n(k) \cdot \frac{(t-1)^n}{n!}$. For all $t \in \mathbb{C}$, we know that $\psi_c(t \cdot x_0, k)$ is a quasi-invariant holomorphic function in k . Hence, the Taylor coefficients $g_n(k) = (d/dt)^n g(t, k)|_{t=1}$ are quasi-invariant holomorphic functions in k . Thus, $g_n(k) \cdot e^{-(x_0 k)} \in Q_{c,x_0}$, for each $n = 0, 1, 2, \dots$. Further, it is known that $\psi_c(0, 0) \neq 0$, see e.g. [EG2]. We deduce that $g_{n_0}(k_0) \neq 0$ for some $n = n_0$, for otherwise $g(t, k_0) = 0$ would imply $g(0, k_0) = \psi_c(0, k_0) = \psi_c(0, 0) = 0$. Set $q_0(k) := g_{n_0}(k) \cdot e^{-(x_0 k)}$. Then for any $p \in Q_{c,x_0}$, we have $p(k) = q_0(k) \cdot f(k)$, where $f(k) \in \mathbb{C}(\mathfrak{h})$ is a rational function regular at $k = k_0$. Observe that f can be written as $\frac{p(k) \cdot e^{(x_0 k)}}{g_{n_0}(k)}$, a ratio of two quasi-invariant functions in k . Hence, $f(k) \in \mathcal{O}_{k_0}(X_c)$ by Lemma 7.3(iii). It follows that $\mathcal{O}_{k_0}(X_c) \otimes_{\mathcal{O}(X_c)} Q_{c,x_0}$ is a free rank 1 module generated by q_0 . $\square$

Definition 7.7. Let $\mathcal{K}_c \subset \mathbb{C}[\mathfrak{h} \times \mathfrak{h}^*]$ be the space of all polynomials P such that the holomorphic function: $(x, k) \mapsto P(x, k)e^{(x, k)}$ is quasi-invariant with respect to both the variable $x \in \mathfrak{h}$ and the variable $k \in \mathfrak{h}^*$, separately.

Observe that the space $\mathcal{K}_c$ has a natural $Q_c \otimes Q_c$ -module structure induced by multiplication in $\mathbb{C}[\mathfrak{h}] \otimes \mathbb{C}[\mathfrak{h}^*]$.

Proposition 7.8. The space $\mathcal{K}_c$ is a projective rank one $Q_c \otimes Q_c$ -module.

Proof. The proof is analogous to the argument above. Specifically, we choose as in Proposition 7.6, a function $g_0 \in \mathcal{K}_c$ that is nonvanishing at a given point (x_0, k_0) . Then, for any $g \in \mathcal{K}_c$, the function $\frac{g}{g_0}$ is a rational function (not involving the exponents) on $\mathfrak{h} \times \mathfrak{h}^*$. Moreover, this function is regular at the point $(x_0, k_0) \in \mathfrak{h} \times \mathfrak{h}^*$ and $W_{x_0} \times W_{k_0}$ -quasi-invariant. Hence, by Lemma 7.3(iii) applied to the Coxeter group $W \times W$ in the vector space $\mathfrak{h} \times \mathfrak{h}^*$, we obtain $\frac{g}{g_0} \in \mathcal{O}_{x_0, k_0}(X_c \times X_c)$. It follows that the function g_0 generates the $\mathcal{O}_{x_0, k_0}(X_c \times X_c)$ -module $\mathcal{O}_{(x_0, k_0)}(X_c \times X_c) \otimes_{Q_c \otimes Q_c} \mathcal{K}_c$. Thus, $\mathcal{K}_c$ is a locally free $Q_c \otimes Q_c$ -module of rank one. $\square$

Corollary 7.9. For any $a \in X_c$, we have: $\mathcal{K}_c|_{X_c \times \{a\}} = Q_{c,a}$.

Proof. We have an obvious evaluation map $\text{ev} : \mathcal{K}_c|_{X_c \times \{a\}} \rightarrow Q_{c,a}$. Both sides are rank one locally free sheaves on $X_c \times \{a\}$. Hence, to prove bijectivity we only need to check that the map is surjective. This reduces to checking that for any x there exists $f \in \mathcal{K}_c|_{X_c \times \{a\}}$ such that $\text{ev}(f)(x)$ is nonzero. But $\text{ev}(f)(x) = f(x, a)$, so one can take $f = g_0$, the same function as has been taken in the proof of Proposition 7.8. $\square$

In terms of the ψ_c -function, the Chalykh-Veselov map $Q_c \rightarrow Q_c^b$, $P \mapsto \mathcal{L}_P$, is defined as follows. Given $P \in Q_c$, let $\mathcal{L}_P \in \mathcal{D}(\mathfrak{h}^{\text{reg}})$ be the unique differential operator such that $(\mathcal{L}_P^{(x)}\psi_c)(x, k) = P(k) \cdot \psi_c(x, k)$, where $u^{(x)}$ stands for the operator u acting in the " x -variable".

Below, we will use the following result due to Chalykh-Veselov (we use the notation $\sigma^{(k)}(p)$ for the leading term of $p \in \mathbb{C}[\mathfrak{h}^{\text{reg}} \times \mathfrak{h}^*]$, viewed as a polynomial in the variable ' k ' only, and write $\sigma(u)$ for the principal symbol of a differential operator u).

Lemma 7.10 ([CV2], p.33). *Let $p \in \mathbb{C}[\mathfrak{h}^{\text{reg}} \times \mathfrak{h}^*]$ be a function in the variables $(x, k) \in \mathfrak{h}^{\text{reg}} \times \mathfrak{h}^*$ such that the function $F : (x, k) \mapsto p(x, k) \cdot e^{(x, k)}$ is quasi-invariant with respect to the variable ' k '. Then*

- (i) *There exists a differential operator $u \in \mathcal{D}(\mathfrak{h}^{\text{reg}})$ such that $F = u^{(x)}\psi_c$.*
- (ii) *We have: $\sigma^{(k)}(p) \in \mathbb{C}[\mathfrak{h}^{\text{reg}} \times \mathfrak{h}^*] \cdot \delta_c(k)$ and, moreover, $\sigma(u) = \frac{\sigma^{(k)}(p)}{\delta_c(x)\delta_c(k)}$.* $\square$

Recall next that in [EG2] we have shown that formula $(f, g)_c := (\mathcal{L}fg)(0)$ gives a non-degenerate symmetric $\mathbb{C}$ -bilinear pairing on the vector space Q_c .

Further, given $P \in Q_c$ we write $\deg^{(x)}P$ for the *degree* of P , viewed as a polynomial in the x -variable only. Also, we write $\text{weight}(u)$ for the weight of a $\mathbb{C}^*$ -homogeneous differential operator $u \in \mathcal{D}(X_c)$ with respect to the $\mathbb{C}^*$ -action and $\text{order}(u)$ for the order of u as a differential operator. We let the ring $Q_c \otimes Q_c^b$ act on $\mathcal{K}_c$ via the identification $Q_c \otimes Q_c \simeq Q_c \otimes Q_c^b$ induced by the Chalykh-Veselov isomorphism $Q_c \xrightarrow{\sim} Q_c^b$ on the second tensor factors.

We define a linear map $K : \mathcal{D}(X_c) \rightarrow \mathcal{K}_c$, given by $u \mapsto K(u) = e^{-(x, k)} \cdot (u^{(x)}\psi_c(x, k))$.

Proposition 7.11. (i) *The map $K : \mathcal{D}(X_c) \rightarrow \mathcal{K}_c$ is a $Q_c \otimes Q_c^b$ -module isomorphism.*

(ii) *For any $u \in \mathcal{D}(X_c)$, we have: $\deg^{(x)}(K(u)) - \deg^{(x)}(\Phi) = \text{weight}(u) + \text{order}(u)$, moreover, this integer is ≥ 0 .*

Proof. First of all, K is obviously injective. So, we need to show that for any $P \in \mathcal{K}_c$, there exists $u \in \mathcal{D}(X_c)$ such that $P = K(u)$. Since $P(x, k)e^{(x, k)}$ is quasi-invariant with respect to k , it follows from Lemma 7.10 that there exists a differential operator $u \in \mathcal{D}(\mathfrak{h}^{\text{reg}})$ such that $(u^{(x)}\psi_c)(x, k) = P(x, k)e^{(x, k)}$.

Further, by [EG2], one has an infinite expansion: $\psi_c(x, k) = \sum \psi_i(k)\psi^i(x)$, where $\psi_i, \psi^i \in Q_c$ are homogeneous polynomials which form dual bases with respect to the nondegenerate form $(-, -)_c$. From the expansion we get: $P(x, k)e^{(x, k)} = (u^{(x)}\psi_c)(x, k) = \sum_i \psi_i(k) \cdot (u^{(x)}\psi^i(x))$. Thus, since $P(x, k)e^{(x, k)}$ is also quasi-invariant with respect to x , we find that $u\psi^i \in Q_c$, $\forall i$. Hence the operator u maps Q_c to Q_c . So, $u \in \mathcal{D}(X_c)$ and $P = K(u)$, as desired. Part (i) is proved.

Now, given $P \in Q_c$ and $u \in \mathcal{D}(X_c)$, we find:

$$\begin{aligned} K(u \circ \mathcal{L}_P)(x, k) \cdot e^{(x, k)} &= (u^{(x)} \circ \mathcal{L}_P^{(x)}\psi_c)(x, k) = u^{(x)}(P(k) \cdot \psi_c(x, k)) \\ &= P(k) \cdot (u^{(x)}\psi_c(x, k)) = P(k) \cdot K(u)(x, k) \cdot e^{(x, k)}. \end{aligned}$$

Compatibility with $Q_c \otimes Q_c^b$ -module structures follows.

Observe further that if u is homogeneous under the $\mathbb{C}^*$ -action (of weight m , say), then $K(u)$ is homogeneous of the same weight. It follows that the highest powers in x always come in

$K(u)$ together with the highest powers in k , i.e. come from the leading term in both k and x simultaneously. Now, the leading term of $K(u)$ is equal to the principal symbol of u times the leading term of Φ . This implies (ii). $\square$

We introduce a *non-standard* increasing filtration on $\mathcal{D}(X_c)$ by

$$\mathcal{D}_j^b(X_c) := \text{span of } \{u \in \mathcal{D}(X_c) \mid \text{weight}(u) + \text{order}(u) \leq j\}, \quad j \in \mathbb{Z}. \quad (7.12)$$

For any homogeneous differential operators u and u' , we have $\text{weight}(u \circ u') = \text{weight}(u) + \text{weight}(u')$, and $\text{order}(u \circ u') \leq \text{order}(u) + \text{order}(u')$. Hence, for any $j, l \in \mathbb{Z}$, we get: $\mathcal{D}_j^b(X_c) \cdot \mathcal{D}_l^b(X_c) \subseteq \mathcal{D}_{j+l}^b(X_c)$. Moreover, since $\text{order}(u \circ u' - u' \circ u) < \text{order}(u) + \text{order}(u')$, we deduce that the associated graded algebra $\text{gr}^b \mathcal{D}(X_c)$ is commutative. Further, by Proposition 7.11(ii), we have: $\mathcal{D}_{-1}^b(X_c) = 0$, so that our filtration is by *non-negative* integers $j \geq 0$. Thus, we have proved the inclusion on the right of (7.13):

$$\mathcal{D}(X_c)_- = \mathcal{D}_0^b(X_c) = Q_c^b, \quad \text{and} \quad [u, \mathcal{D}_l^b(X_c)] \subset \mathcal{D}_{j+l-1}^b(X_c), \quad \forall u \in \mathcal{D}_j^b(X_c). \quad (7.13)$$

To prove equations on the left of (7.13), let $u \in \mathcal{D}(X_c)$ be a homogeneous differential operator such that $\text{weight}(u) + \text{order}(u) = 0$, and let $K(u)$ be its kernel. Since $K(u)$ is quasi-invariant with respect to the ' x '-variable, Lemma 7.10(i) implies that there exists a differential operator $p \in \mathcal{D}(\mathfrak{h}^{\text{reg}})$ (in the variable $k \in \mathfrak{h}^{\text{reg}}$) such that $K(u) = p^{(k)}\psi_c$. Now, the proof of Proposition 7.11(ii) shows that $\text{weight}(u) + \text{order}(u) = 0$ if and only if $\deg^{(x)}(p^{(k)}\psi_c) = \deg^{(x)}\Phi$. If the latter equation holds, then Lemma 7.10(ii) forces the differential operator $p \in \mathcal{D}(\mathfrak{h}^{\text{reg}})$ to have order zero, i.e., $p = p(k) \in \mathbb{C}[k] = \mathbb{C}[\mathfrak{h}^*]$ is a multiplication operator. Hence, $K(u) \cdot e^{(x,k)} = (p^{(k)}\psi_c)(x, k) = p(k) \cdot \psi_c(x, k) = p(k) \cdot \Phi(x, k) \cdot e^{(x,k)}$. Thus, we have proved that $\text{weight}(u) + \text{order}(u) = 0$ holds if and only if $K(u)$ has the form $p \cdot \Phi$, where p is a polynomial in the variable ' k ' only. But then we must have $p \in Q_c$ and $u = \mathcal{L}_p$. This completes the proof of (7.13). $\square$

Next, identify $\mathfrak{h}^*$ with $\mathfrak{h}$ via an invariant form, and write $P^b \in \mathbb{C}[\mathfrak{h} \times \mathfrak{h}^*]$ for the polynomial obtained from P by the flip of variables: $x \leftrightarrow k$. It is clear that $P^b \in \mathcal{K}_c$. We write $u^b := K^{-1}(P^b) \in \mathcal{D}(X_c)$ for the differential operator corresponding to the kernel P^b .

Given a linear map $u : Q_c \rightarrow Q_c$, let $u^\dagger : Q_c \rightarrow Q_c$ denote the adjoint map with respect to the nondegenerate form $(-, -)_c$ on Q_c .

Lemma 7.14. (i) $(u^b(f), g)_c = (f, u(g))_c$, $\forall f, g \in Q_c$ and $u \in \mathcal{D}(X_c)$. That is, $u^b = u^\dagger$, and the assignment: $u \mapsto u^\dagger = u^b$ gives a well-defined algebra anti-involution on $\mathcal{D}(X_c)$.

(ii) We have: $\mathcal{L}_p = P^b$, for any $P \in Q_c$.

Proof. By the results of [EG2], it is sufficient to prove the identity $(u^b(f), g)_c = (f, u(g))_c$ for $f(x) = \psi_c(x, a)$, $\forall a \in \mathfrak{h}^*$. Recall that $(u^{(x)})^b \psi_c(x, a) = (u^{(k)}\psi_c(x, k))|_{k=a}$. Thus $(u^b(f), g)_c = (ug)(a) = (f, u(g))_c$, and the first equation of part (i) follows. This yields $u^b = u^\dagger$, for any $u \in \mathcal{D}(X_c)$. Thus, we have proved that $u \in \mathcal{D}(X_c) \implies u^\dagger \in \mathcal{D}(X_c)$. Since $u^\dagger \circ v^\dagger = (v \circ u)^\dagger$, part (i) follows. Part (ii) follows from the formula $(\mathcal{L}_p)^\dagger = p$, $\forall p \in Q_c$, proved in [EG2]. $\square$

Proof of Theorem 7.5. We prove first that Q_c^b is the centralizer of the Calogero-Moser operator in $\mathcal{D}(\mathfrak{h}^{\text{reg}})_-$. To this end, write $\sigma(u) \in \mathbb{C}(\mathfrak{h})[\mathfrak{h}^*]$ for the principal symbol of a differential operator $u \in \mathcal{D}(\mathbb{K})$. Observe that, for any $p \in Q_c$, we have: $\sigma(\mathcal{L}_p) = p$. Now, let $u \in \mathcal{D}(\mathfrak{h}^{\text{reg}})_-$ be a homogeneous operator such that $[u, \mathcal{L}_c] = 0$. We will prove that $u \in Q_c^b$. It suffices to show that, $\sigma(u) = p \in Q_c$. Then, the operator $u' = u - \mathcal{L}_p$ commutes with $\mathcal{L}_c$. Since $\text{order}(u') < \text{order}(u)$ and $\text{weight}(u') = \text{weight}(u)$, we deduce that $\text{weight}(u') + \text{order}(u') < 0$. Thus, $u' = 0$ by (7.13), and $u = \mathcal{L}_p$.

Observe further that the leading term of u must have constant coefficients (i.e., for the principal symbol we have: $\sigma(u) \in \mathbb{C}[\mathfrak{h}^*]$) since $\text{weight}(u) + \text{order}(u) \leq 0$. Hence $\sigma(u) = p$, for some polynomial $p \in \mathbb{C}[\mathfrak{h}]$. Also, we claim that u commutes with Q_c^b . Indeed, if $u' \in Q_c^b$ then for the operator $[u, u']$, the sum of order and degree is strictly negative, so the operator must be zero. Thus, u preserves each of the weight-spaces of Q_c^b in the vector space $\bigoplus_{k \in \mathbb{C}} \mathbb{C}[\mathfrak{h}^{\text{reg}}] \cdot e^{(k, x)}$. These weight spaces are known, cf. e.g. [CV], to be 1-dimensional and spanned by the function $\psi_c(-, k)$. It follows that $u^{(x)}\psi_c(x, k) = p(k)\psi_c(x, k)$. Therefore, p is quasi-invariant. Thus, we have proved that the centralizer of L_c in $\mathcal{D}(\mathfrak{h}^{\text{reg}})_-$ equals Q_c^b .

All the remaining claims of part (i) of the Theorem now follow from (7.13). Part (ii) follows from Proposition 7.11(i) and Proposition 7.8. $\square$

Remark. The adjoint action on $\mathcal{D}(X_c)$ of the subalgebra $Q_c = \mathbb{C}[X_c]$ is locally nilpotent by the definition of differential operators. Since $u \mapsto u^\dagger$ is an anti-homomorphism, Lemma 7.14(i) implies that the adjoint action on $\mathcal{D}(X_c)$ of the subalgebra Q_c^b is locally nilpotent. This yields formula (6.5) in full generality.

An alternative proof of nilpotency of the adjoint Q_c^b -action on $\mathcal{D}(X_c)$ follows from formulas (7.13), which yield: $[u, \mathcal{D}_l^b(X_c)] \subset \mathcal{D}_{l-1}^b(X_c)$, for any $u \in Q_c^b$ and $l \geq 0$. $\square$

Let $\mathcal{D}_c$ denote the subalgebra in $\mathcal{D}(X_c)$ generated by $Q_c = \mathbb{C}[X_c]$ and Q_c^b . In the previous section we have introduced the subalgebra $\mathcal{B}_c$ generated by Q_c^W and $(Q_c^b)^W$. Thus we have

$$\mathbf{e}H_c\mathbf{e} \simeq \mathcal{B}_c \subset \mathcal{D}_c \subset \mathcal{D}(X_c). \quad (7.15)$$

The inclusions in (7.15) give an algebra imbedding: $\mathbf{e}H_c\mathbf{e} \hookrightarrow \mathcal{D}(X_c)$, making $\mathcal{D}(X_c)$ an $\mathbf{e}H_c\mathbf{e}$ -bimodule.

Proposition 7.16. (i) $\mathcal{D}(X_c)$ is a Harish-Chandra $\mathbf{e}H_c\mathbf{e}$ -bimodule.

(ii) The group $SL_2(\mathbb{C})$ acts on this Harish-Chandra bimodule by algebra automorphisms; the Fourier transform interchanges Q_c and Q_c^b .

Proof. To show that $\mathcal{D}(X_c)$ is a Harish-Chandra $\mathbf{e}H_c\mathbf{e}$ -bimodule, recall that we have proved that $\mathcal{D}(X_c) \simeq \mathcal{K}_c$, as a $\mathbb{C}[\mathfrak{h}]^W \otimes \mathbb{C}[\mathfrak{h}^*]^W$ -module. Further, $\mathcal{K}_c$, being a submodule in $\mathbb{C}[\mathfrak{h} \times \mathfrak{h}^*]$, is clearly a finitely generated $\mathbb{C}[\mathfrak{h}]^W \otimes \mathbb{C}[\mathfrak{h}^*]^W$ -module. It follows that $\mathcal{D}(X_c)$ is a finitely generated $\mathbf{e}H_c\mathbf{e} \otimes (\mathbf{e}H_c\mathbf{e})^{\text{op}}$ -module. By a remark above (7.15) we know that the adjoint action on $\mathcal{D}(X_c)$ of any element of either $\mathbb{C}[\mathfrak{h}]^W \subset Q_c$ or $\mathbb{C}[\mathfrak{h}^*]^W \subset Q_c^b$ is locally nilpotent. This proves (i).

To prove (ii) note that the Fourier automorphism on $\mathbf{e}H_c\mathbf{e}$ interchanges $\mathbb{C}[\mathfrak{h}]^W\mathbf{e}$ with $\mathbb{C}[\mathfrak{h}^*]^W\mathbf{e}$, see [EG]. But Q_c is obviously the centralizer of $\mathbb{C}[\mathfrak{h}]^W\mathbf{e}$ in $\mathcal{D}(X_c)$. Thus, the image of Q_c under the Fourier automorphism equals the centralizer of $\mathbb{C}[\mathfrak{h}^*]^W\mathbf{e}$, which is exactly Q_c^b . $\square$

Further, observe that the natural W -action on X_c induces a W -action on $\mathcal{D}(X_c)$ by algebra automorphisms. This W -action clearly commutes with both the left and right action of $\mathbf{e}H_c\mathbf{e}$ on $\mathcal{D}(X_c)$. Thus, we have a direct sum decomposition

$$\mathcal{D}(X_c) = \bigoplus_{\tau \in \text{Irrep}(W)} \tau^* \otimes D(\tau), \quad \text{where } D(\tau) = (\tau \otimes \mathcal{D}(X_c))^W \text{ are } \mathbf{e}H_c\mathbf{e}\text{-bimodules.}$$

Note that, for any $\tau \in \text{Irrep}(W)$, we have: $D(\tau) \neq 0$, since $0 \neq (\tau \otimes Q_c)^W \subset D(\tau)$.

Theorem 7.17. The bimodules $\{D(\tau)\}_{\tau \in \text{Irrep}(W)}$, are pairwise non-isomorphic simple Harish-Chandra $\mathbf{e}H_c\mathbf{e}$ -bimodules, for any $c \in \mathbb{Z}[R]_+^W$.

We start the proof with the following lemma, which is perhaps well known.

Lemma 7.18. *Let X be a smooth affine algebraic variety over $\mathbb{C}$ with a free action of a finite group Γ . Let τ be an irreducible Γ -module. Then the space $\mathcal{D}_\tau(X) := \text{Hom}_\Gamma(\tau, \mathcal{D}(X))$ is a simple bimodule over $\mathcal{D}(X/\Gamma)$.*

Proof. Using the equivalence between left and right $\mathcal{D}$ -modules, we can regard $\mathcal{D}_\tau(X)$ as a $\mathcal{D}$ -module on $X/\Gamma \times X/\Gamma$. It is concentrated on the diagonal, and is the direct image of the irreducible local system on the diagonal, corresponding to the representation τ of Γ . By Kashiwara's theorem, the direct image functor is an equivalence between the category of $\mathcal{D}$ -modules on the subvariety and $\mathcal{D}$ -modules on the ambient variety supported on the subvariety. Thus, $\mathcal{D}_\tau(X)$ is an irreducible $\mathcal{D}$ -module, as desired. $\square$

Proof of Theorem 7.17. We know that $\mathcal{D}(X_c)$, hence any its W -isotypic component $D(\tau)$, is a Harish-Chandra $\mathbf{eH}_c\mathbf{e}$ -bimodule. We first prove simplicity of the bimodule $D(\tau)$, which is parallel to the proof of simplicity of the algebra $\mathbf{H}_c$. Consider the localization $D(\tau)|_{\mathfrak{h}^{\text{reg}}/W}$ of $D(\tau)$. This is a bimodule over $(\mathbf{eH}_c\mathbf{e})|_{\mathfrak{h}^{\text{reg}}/W} = \mathcal{D}(\mathfrak{h}^{\text{reg}})^W$, which is isomorphic to $(\tau \otimes \mathcal{D}(\mathfrak{h}^{\text{reg}}))^W$.

Let J be a nonzero subbimodule of $D(\tau)$. Consider the Harish-Chandra $\mathbf{eH}_c\mathbf{e}$ -bimodule $V = D(\tau)/J$, and its localization $V|_{\mathfrak{h}^{\text{reg}}/W}$. By Lemma 7.18, $D(\tau)|_{\mathfrak{h}^{\text{reg}}/W}$ is a simple $(\mathbf{eH}_c\mathbf{e})|_{\mathfrak{h}^{\text{reg}}/W}$ -module. Hence, $V|_{\mathfrak{h}^{\text{reg}}/W} = (D(\tau)|_{\mathfrak{h}^{\text{reg}}/W})/(J|_{\mathfrak{h}^{\text{reg}}/W}) = 0$. Corollary 4.5 implies that $V = 0$. Hence $J = D(\tau)$, and $D(\tau)$ is a simple $\mathbf{eH}_c\mathbf{e}$ -bimodule.

Finally, the bimodules $D(\tau)$, $\tau \in \text{Irrep}(W)$, are pairwise nonisomorphic since the localized bimodules $D(\tau)|_{\mathfrak{h}^{\text{reg}}/W} = (\tau \otimes \mathcal{D}(\mathfrak{h}^{\text{reg}}))^W$ are already nonisomorphic. The latter is clear since these modules, viewed as $\mathcal{D}$ -modules on $\mathfrak{h}^{\text{reg}} \times \mathfrak{h}^{\text{reg}}$ give rise to pairwise non-isomorphic local systems on the diagonal in $\mathfrak{h}^{\text{reg}}/W \times \mathfrak{h}^{\text{reg}}/W$. $\square$

The following is an analogue of the Levasseur-Stafford theorem

Theorem 7.19. *For any $c \in \mathbb{Z}[R]_+^W$, the algebra $\mathcal{D}(X_c)$ is generated by the two commutative subalgebras, Q_c and Q_c^b .*

Proof. Recall, see (7.15), that $\mathcal{D}_c$ denotes the subalgebra of $\mathcal{D}(X_c)$ generated by Q_c and Q_c^b . Since $\mathbf{eH}_c\mathbf{e}$ is generated by $\mathbb{C}[\mathfrak{h}]^W\mathbf{e}$ and $\mathbb{C}[\mathfrak{h}^*]^W\mathbf{e}$, the subalgebra $\mathcal{D}_c$ is a $\mathbb{C}W \otimes \mathbf{eH}_c\mathbf{e} \otimes (\mathbf{eH}_c\mathbf{e})^{op}$ -submodule in $\mathcal{D}(X_c)$. The $\mathbb{C}W \otimes \mathbf{eH}_c\mathbf{e} \otimes (\mathbf{eH}_c\mathbf{e})^{op}$ -module decomposition: $\mathcal{D}(X_c) = \bigoplus_{\tau \in \text{Irrep}(W)} \tau^* \otimes D(\tau)$, and Theorem 7.17 imply that $\mathcal{D}_c = \bigoplus_{\tau \in S} \tau^* \otimes D(\tau)$, where summation runs over a certain set $S \subseteq \text{Irrep}(W)$ of irreducible W -modules. However, $\mathcal{D}_c$ contains Q_c , hence every irreducible W -module occurs in $\mathcal{D}_c$ with nonzero multiplicity. Thus, the set S has to be the entire set $\text{Irrep}(W)$, and we are done. $\square$

Theorem 7.20. *The algebra $\mathcal{D}(X_c)$ is simple, for any $c \in \mathbb{Z}[R]_+^W$.*

Proof. We begin with the following general

Claim. *Let X be an irreducible affine variety, and let $A \subset \mathbb{C}[X]$ be a subalgebra such that $\mathbb{C}[X]$ is a finite module over A . Then any nonzero two-sided ideal of $\mathcal{D}(X)$ intersects A non-trivially.*

To prove the Claim, let J be a non-trivial two-sided ideal of $\mathcal{D}(X)$. Choose a nonzero differential operator, say u , in J of the minimal order. Since $[u, f] \in J$ and $\text{order}([u, f]) < \text{order}(u)$, for any $f \in \mathbb{C}[X]$, we conclude that $[u, f] = 0$ and hence $u \in \mathbb{C}[X]$. It follows that $u \in J \cap \mathbb{C}[X]$. Now, being finite as a module over A , the ring $\mathbb{C}[X]$ is integral over A . Hence, u satisfies an equation $u^n + a_1 \cdot u^{n-1} + \dots + a_{n-1} \cdot u + a_0 = 0$, with coefficients in A . In particular, there is a nonzero $g \in \mathbb{C}[X]$ such that $g \cdot u \in A$. Since $u \in J \implies g \cdot u \in J$, we have $J \cap A \neq 0$, and the Claim follows.

To prove the Theorem, assume $J \subseteq \mathcal{D}(X_c)$ is a nonzero two-sided ideal. The Claim above applied to $A = \mathbb{C}[\mathfrak{h}]^W \subset \mathbb{C}[X_c]$, yields $\mathbb{C}[\mathfrak{h}]^W \cap J \neq 0$. Thus, J nontrivially intersects $\mathbf{eH}_c\mathbf{e}$. But $\mathbf{eH}_c\mathbf{e}$ is simple, so $J \cap \mathbf{eH}_c\mathbf{e} = \mathbf{eH}_c\mathbf{e}$. Hence $1 \in J$, and $J = \mathcal{D}(X_c)$. $\square$

The following corollary of Theorem 7.20 has been first proved in [EG2] by a different method.

Corollary 7.21. *For any $c \in \mathbb{Z}_+(R)^W$, the variety X_c is Cohen-Macaulay.*

Proof. Follows from a theorem of Van den Bergh (see [VdB], Theorem 6.2.5) stating that X is a Cohen-Macaulay variety whenever the ring $\mathcal{D}(X)$ is simple. $\square$

Remark. X_c is Cohen-Macaulay $\iff Q_c = \mathbb{C}[X_c]$ is a free $\mathbb{C}[\mathfrak{h}]^W$ -module. The freeness of Q_c over $\mathbb{C}[\mathfrak{h}]^W$ has been conjectured by Feigin-Veselov and proved in [FV] for dihedral groups, and in [EG2] in general.

Proposition 7.22. *The imbedding: $\mathbf{eH}_c\mathbf{e} \hookrightarrow \mathcal{D}(X_c)^W$ is an algebra isomorphism.*

Proof. Let $u \in \mathcal{D}(X_c)$ be a homogeneous operator and $K_0(u)$ be the highest term of the corresponding kernel $K(u) \in \mathcal{K}_c$, cf. Definition 7.7. It is clear that $K_0(u)(x, k) = \delta_c(x)\delta_c(k) \cdot \text{Symbol}(u)$. The principal symbol is always polynomial in k , so $K_0(u)(x, k)$ is always divisible by $\delta_c(k)$. But by symmetry of $x \longleftrightarrow k$, it is also always divisible by $\delta_c(x)$. Thus, we have proved that the principal symbol of any differential operator $u \in \mathcal{D}(X_c)$ is polynomial, and hence $\text{gr}(\mathcal{D}(X_c)^W) \subset \mathbb{C}[\mathfrak{h}^* \times \mathfrak{h}]^W$.

On the other hand, $\text{gr}(\mathbf{eH}_c\mathbf{e}) = \mathbb{C}[\mathfrak{h}^* \times \mathfrak{h}]^W$. Hence, the associated graded map: $\text{gr}(\mathbf{eH}_c\mathbf{e}) \rightarrow \text{gr}(\mathcal{D}(X_c)^W)$ is surjective. The Proposition follows. $\square$

8 Translation functors and Morita equivalence

Theorem 8.1. *Let $c \in \mathbb{C}[R]_{\text{reg}}^W$. Then, for any $m \in \mathbb{Z}[R]^W$, the algebras $\mathbf{H}_c$ and $\mathbf{H}_{c-m}$ (resp. the algebras $\mathbf{eH}_c\mathbf{e}$ and $\mathbf{eH}_{c-m}\mathbf{e}$) are Morita equivalent.*

Proof. We may assume without loss of generality that $m \in \mathbb{Z}[R]_+^W$. Recall that the algebra $\mathbf{H}_c$ is simple for any $c \in \mathbb{C}[R]_{\text{reg}}^W$, therefore both $\mathbf{H}_c$ and $\mathbf{H}_{c-m}$ are simple. Thus, using Proposition 4.11 and Corollary 4.2 we deduce

$$\mathbf{H}_c \xrightarrow[\text{Lemma 4.1}]{\text{Morita equivalence}} \mathbf{e}_\varepsilon \mathbf{H}_c \mathbf{e}_\varepsilon \xrightarrow[\text{Prop. 4.11}]{\text{Prop. 4.11}} \mathbf{eH}_{c-1_\varepsilon} \mathbf{e} \xrightarrow[\text{equivalence}]{\text{Morita}} \mathbf{H}_{c-1_\varepsilon}.$$

Recall further that the functions of the form 1_ε , where $\varepsilon : W \rightarrow \{\pm 1\}$ is a multiplicative character, see above Proposition 4.11, are known to generate $\mathbb{Z}[R]_+^W$ as a semi-group. Thus m is a sum of functions of type 1_ε , and an easy induction on the number of summands completes the proof. $\square$

Lemma 8.2. *For any $c \in \mathbb{Z}[R]^W$, the algebra $\mathbf{H}_c$ is Morita equivalent to $\mathcal{D}(\mathfrak{h})\#W$.*

Proof. For $c \in \mathbb{Z}[R]^W$, Theorem 8.1 yields Morita equivalence of the algebras $\mathbf{H}_c$ and $\mathbf{H}_0$. But $\mathbf{H}_0 \simeq \mathcal{D}(\mathfrak{h})\#W$, and the result follows. $\square$

Remark. If $\mathfrak{h} = \mathbb{C}$ and $W = \mathbb{Z}/2\mathbb{Z}$, then $\mathbf{eH}_c\mathbf{e}$ is a quotient of $U(\mathfrak{sl}_2)$, and Theorem 8.1 is known and due to Stafford [St]. $\square$

By analogy with Lie theory, we denote by $\mathcal{HC}_c(\mathbf{H})$ the category of Harish-Chandra $\mathbf{H}_c$ -bimodules. In a similar way, we may consider the category $\mathcal{HC}_c(\mathbf{eHe})$ of Harish-Chandra bimodules over the algebra $\mathbf{eHe}$. Recall that the category of bimodules over any algebra A has a monoidal structure given by the tensor product of bimodules over A .

Lemma 8.3. *The tensor product of two Harish-Chandra bimodules is again a Harish-Chandra bimodule.*

Proof. Let $V, V' \in \mathcal{HC}_c(\mathbf{H})$. The Leibniz rule for derivations shows that the adjoint action on $V \otimes_{\mathbf{H}_c} V'$ of any element of $\mathbb{C}[\mathfrak{h}]^W$ and of any element of $\mathbb{C}[\mathfrak{h}^*]^W$ is locally nilpotent. Thus, it suffices to prove that $V \otimes_{\mathbf{H}_c} V'$ is a finitely generated $\mathbf{H}_c \otimes \mathbf{H}_c^{op}$ -module.

To this end, choose a finite set $\{v_i\}$, resp. $\{v'_j\}$, of generators of V , resp. of V' , as a $\mathbb{C}[\mathfrak{h}]^W$ - $\mathbb{C}[\mathfrak{h}^*]^W$ -bimodule, see Lemma 3.3(ii). Then we get

$$V \otimes_{\mathbf{H}_c} V' = \sum_{i,j} (\mathbb{C}[\mathfrak{h}]^W \cdot v_i \cdot \mathbb{C}[\mathfrak{h}^*]^W) \otimes_{\mathbf{H}_c} (\mathbb{C}[\mathfrak{h}]^W \cdot v'_j \cdot \mathbb{C}[\mathfrak{h}^*]^W). \quad (8.4)$$

Now Lemma 2.8 insures, due to nilpotency of the adjoint action, that the subspace $\mathbb{C}[\mathfrak{h}]^W \cdot v \cdot \mathbb{C}[\mathfrak{h}]^W \subset V$ is finitely generated as a left $\mathbb{C}[\mathfrak{h}]$ -module. Similarly, $\mathbb{C}[\mathfrak{h}^*]^W \cdot v' \cdot \mathbb{C}[\mathfrak{h}^*]^W$ is finitely generated as a right $\mathbb{C}[\mathfrak{h}^*]^W$ -module, for any $v' \in V'$. We conclude that there are finite sets $\{u_{ik}\} \subset V$ and $\{u'_{jl}\} \subset V'$, such that: $\mathbb{C}[\mathfrak{h}]^W \cdot v_i \cdot \mathbb{C}[\mathfrak{h}]^W \subset \sum_k \mathbb{C}[\mathfrak{h}]^W \cdot u_{ik}$, and $\mathbb{C}[\mathfrak{h}^*]^W \cdot v'_j \cdot \mathbb{C}[\mathfrak{h}^*]^W \subset \sum_l \mathbb{C}[\mathfrak{h}^*]^W \cdot u'_{jl}$.

Further, by the Poincaré-Birkhoff-Witt isomorphism (1.2), there exists a finite dimensional subspace $E \subset \mathbf{H}_c$ such that $\mathbb{C}[\mathfrak{h}^*]^W \otimes_{\mathbf{H}_c} \mathbb{C}[\mathfrak{h}]^W \subset \mathbb{C}[\mathfrak{h}]^W \cdot E \cdot \mathbb{C}[\mathfrak{h}^*]^W$ (inclusion of subspaces in $\mathbf{H}_c \otimes_{\mathbf{H}_c} \mathbf{H}_c$). The proof is now completed by the following inclusions

$$\begin{aligned} \text{RHS of (8.4)} &\subseteq \sum_{i,j} \mathbb{C}[\mathfrak{h}]^W \cdot v_i \cdot \mathbb{C}[\mathfrak{h}]^W \cdot E \cdot \mathbb{C}[\mathfrak{h}^*]^W \cdot v'_j \cdot \mathbb{C}[\mathfrak{h}^*]^W \\ &\subseteq \sum_{i,j,k,l} \mathbb{C}[\mathfrak{h}]^W \cdot u_{ik} \cdot E \cdot u'_{jl} \cdot \mathbb{C}[\mathfrak{h}^*]^W \subseteq \sum_{i,j,k,l} \mathbf{H}_c \cdot u_{ik} \cdot E \cdot u'_{jl} \cdot \mathbf{H}_c. \quad \square \end{aligned}$$

Thus, for each $c \in \mathbb{C}[R]^W$, the category $\mathcal{HC}_c(\mathbf{H})$, resp. $\mathcal{HC}_c(\mathbf{eHe})$, has a monoidal structure induced by the tensor product of bimodules. In case of $c \in \mathbb{Z}[R]^W_+$ a complete structure of these monoidal categories can be described as follows. First, recall that for any finite dimensional (not necessarily irreducible) W -representation τ , the space $D(\tau) := (\tau \otimes \mathcal{D}(X_c))^W$ is a Harish-Chandra $\mathbf{eHe}$ -bimodule, cf. Theorem 7.17. Let $\text{Rep}(W)$ denote the tensor category of finite dimensional representations of the group W .

Theorem 8.5. *For any $c \in \mathbb{Z}[R]^W_+$, the assignment $\tau \mapsto D(\tau) := (\tau \otimes \mathcal{D}(X_c))^W$ gives an equivalence $\Xi : \text{Rep}(W) \xrightarrow{\sim} \mathcal{HC}_c(\mathbf{eHe})$ of monoidal categories. The monoidal structure on the functor Ξ is given, for any $\tau, \sigma \in \text{Irrep}(W)$, by a canonical $\mathbf{eHe}$ -bimodule isomorphism: $D(\tau) \otimes_{\mathbf{eHe}} D(\sigma) \xrightarrow{\sim} D(\tau \otimes \sigma)$ induced by multiplication in the algebra $\mathcal{D}(X_c)$:*

$$(\tau \otimes \mathcal{D}(X_c))^W \bigotimes_{\mathbf{eHe}} (\sigma \otimes \mathcal{D}(X_c))^W \xrightarrow{\sim} ((\tau \otimes \sigma) \otimes \mathcal{D}(X_c))^W.$$

Similarly, there is an equivalence of monoidal categories $\text{Rep}(W) \xrightarrow{\sim} \mathcal{HC}_c(\mathbf{H})$.

Proof. We first consider the case $c = 0$. In this case we have $\mathbf{H}_0 \simeq \mathcal{D}(\mathfrak{h}) \# W$ and $\mathbf{eHe} \simeq \mathcal{D}(\mathfrak{h})^W$.

Since $\mathbf{H}_0 = \mathcal{D}(\mathfrak{h}) \# W$, a Harish-Chandra $\mathbf{H}_0$ -bimodule is in particular a $\mathcal{D}(\mathfrak{h})$ -bimodule with a $W \times W$ -action, that is a $W \times W$ -equivariant $\mathcal{D}$ -module, V , on $\mathfrak{h} \times \mathfrak{h}$. The bound given in (3.4) shows that $\text{Ch}V$, the *characteristic variety* of V (as a $\mathcal{D}(\mathfrak{h} \times \mathfrak{h})$ -module), is contained

in $\left(\bigcup_{w \in W} \text{Graph}(w)\right) \times \left(\bigcup_{y \in W} \text{Graph}(y)\right)$, which is a subvariety in $T^*(\mathfrak{h} \times \mathfrak{h})$ of dimension $\frac{1}{2} \dim T^*(\mathfrak{h} \times \mathfrak{h})$. Hence, V is a holonomic $\mathcal{D}$ -module. Moreover, since $\text{Ch } V$ must be a Lagrangian subvariety, we deduce

$$\text{Ch } V \subseteq \bigcup_{w \in W} T_{\text{Graph}(w)}^*(\mathfrak{h} \times \mathfrak{h}). \quad (8.6)$$

Given $w \in W$, let $i_w : \text{Graph}(w) \hookrightarrow \mathfrak{h} \times \mathfrak{h}$ denote the imbedding. Kashiwara's theorem and estimate (8.6) imply that every irreducible subquotient of V , viewed as a $\mathcal{D}$ -module, must be of the form $(i_w)_* E$, where E is an irreducible local system, i.e., a vector bundle on $\text{Graph}(w)$ with flat connection. The condition of ad -nilpotency of the action on $(i_w)_* E \subset V$ of the algebra $\mathbb{C}[\mathfrak{h}^*]^W$ forces the connection to be trivial. We deduce that all irreducible subquotients of V are of the form $(i_w)_* \mathbb{C}[\text{Graph}(w)]$. Semisimplicity of the category $\mathcal{HC}_0(\mathbf{H})$ follows from this, since it is well-known that there are no extensions between $\mathcal{D}$ -modules of the form above. We conclude that the category $\mathcal{HC}_0(\mathbf{H})$ is semisimple.

We have shown that any object $V \in \mathcal{HC}_0(\mathbf{H})$ can be uniquely written, viewed as a $\mathcal{D}(\mathfrak{h} \times \mathfrak{h})$ -module, in the form $V = \bigoplus_{w \in W} V_w \otimes_{\mathbb{C}} (i_w)_* \mathbb{C}[\text{Graph}(w)]$, where V_w are finite dimensional vector spaces. Moreover, the left and right actions on V of the subalgebra $\mathbb{C}W \subset \mathbf{H}_0$ induce, for any $y \in W$, the natural isomorphisms $L_y : V_w \xrightarrow{\sim} V_{yw}$ and $R_y : V_w \xrightarrow{\sim} V_{wy}$, which are multiplicative, and commute: $R_y L_z = L_z R_y$, $\forall y, z \in W$. This means that the vector space $V_W := \bigoplus_{w \in W} V_w$ is a representation of $W \times W$, via $(y_1, y_2)v = L_{y_1} R_{y_2}^{-1} v$. It is clear that $V_W = \text{Ind}_{W_{\text{diag}}}^{W \times W} V_1$, where the action on V_1 of the diagonal subgroup $W_{\text{diag}} \subset W \times W$ is given by $w : v \mapsto L_w R_w^{-1}(v)$, as above. Moreover, we see that $V \simeq \text{Ind}_A^{\mathbf{H}_0 \otimes \mathbf{H}_0^{op}} \left((i_{\text{diag}})_* (V_1 \otimes \mathbb{C}[\mathfrak{h}_{\text{diag}}]) \right)$, where A is the preimage of $\mathbb{C}[W_{\text{diag}}]$ under the natural projection $\mathbf{H}_0 \otimes \mathbf{H}_0^{op} = \mathcal{D}(\mathfrak{h} \times \mathfrak{h}) \# (W \times W) \rightarrow \mathbb{C}[W \times W]$. Here $V_1 \otimes \mathbb{C}[\mathfrak{h}_{\text{diag}}]$ is a W -equivariant $\mathcal{D}(\mathfrak{h})$ -module, and $i_{\text{diag}} : \mathfrak{h}_{\text{diag}} \hookrightarrow \mathfrak{h} \times \mathfrak{h}$ is the diagonal map.

Next, recall the following general result on Morita equivalence. Let A be an algebra and $\mathbf{e} \in A$ an idempotent, such that $\mathbf{e}A\mathbf{e} = A$. Then, *the natural Morita equivalence between A -bimodules and $\mathbf{e}A\mathbf{e}$ -bimodules is an equivalence of monoidal categories.*

In the special case $A = \mathbf{H}_c$ the result above implies that the categories $\mathcal{HC}_0(\mathbf{H})$ and $\mathcal{HC}_0(\mathbf{eH}_c\mathbf{e})$ are equivalent as monoidal categories, due to Morita equivalence of the algebras $\mathcal{D}(\mathfrak{h}) \# W$ and $\mathcal{D}(\mathfrak{h})^W$ proved in [Mo]. In particular, the category $\mathcal{HC}_0(\mathbf{eH}_c\mathbf{e})$ is also semisimple.

Further it is easy to see, e.g. ([LS2], §§2-3) or Theorem 7.17 (in the special case $c = 0$) that, for each $\tau \in \text{Irrep}(W)$, the W -isotypic component $(\tau \otimes_{\mathbb{C}} \mathcal{D}(\mathfrak{h}))^W$ is a simple $\mathbf{eH}_c\mathbf{e}$ -bimodule. Moreover, any simple object of $\mathcal{HC}_0(\mathbf{eH}_c\mathbf{e})$ is isomorphic, by Morita equivalence, to a bimodule of this type. Thus, simple objects of the category $\mathcal{HC}_0(\mathbf{eH}_c\mathbf{e})$ are parametrized by the set $\text{Irrep}(W)$. This completes the proof of the Theorem for $c = 0$.

In the general case of an arbitrary $c \in \mathbb{Z}[R]_+^W$, the result follows from that for $c = 0$ via Morita equivalence of Proposition 8.11 below (which is independent of the intervening material). $\square$

Corollary 8.7. *Both $\mathcal{HC}_c(\mathbf{eH}_c\mathbf{e})$ and $\mathcal{HC}_c(\mathbf{H})$ are semisimple tensor categories.* $\square$

To proceed further, it will be useful to generalize the setup and to consider bimodules over two Cherednik algebras with possibly *different* parameters. Thus, we fix $c, c' \in \mathbb{C}[R]^W$, and let $\mathbf{H}_c, \mathbf{H}_{c'}$ be the corresponding Cherednik algebras. There are canonical imbeddings of both $\mathbb{C}[\mathfrak{h}]^W$ and $\mathbb{C}[\mathfrak{h}^*]^W$ into any of the two algebras $\mathbf{H}_c$ or $\mathbf{H}_{c'}$. We have, cf. Definition 3.2

Definition 8.8. A finitely generated $\mathbf{H}_c \otimes \mathbf{H}_{c'}^{op}$ -module V is called a Harish-Chandra $(\mathbf{H}_c, \mathbf{H}_{c'})$ -bimodule if, for any $a \in \mathbb{C}[\mathfrak{h}]^W$ and any $a \in \mathbb{C}[\mathfrak{h}^*]^W$, the $\text{ad } a$ -action on V is locally nilpotent.

We write $\mathcal{HC}_{c,c'}(\mathbf{H})$, resp. $\mathcal{HC}_{c,c'}(\mathbf{eHe})$, for the category of Harish-Chandra $(\mathbf{H}_c\text{-}\mathbf{H}_{c'})$ -bimodules, resp. $(\mathbf{eH}_c\mathbf{e}\text{-}\mathbf{eH}_{c'}\mathbf{e})$ -bimodules. Most of the results on Harish-Chandra bimodules proved in §§3-4 extend verbatim to this ‘two-parameter’ setting.

Let V be an $(\mathbf{H}_c\text{-}\mathbf{H}_{c'})$ -bimodule, resp. $(\mathbf{eH}_c\mathbf{e}\text{-}\mathbf{eH}_{c'}\mathbf{e})$ -bimodule. We say that a vector $v \in V$ is of order $\leq d$ if for any $f \in \mathbb{C}[\mathfrak{h}]^W$ and $g \in \mathbb{C}[\mathfrak{h}^*]^W$, one has $(\text{ad } f)^{d_1+1}v = (\text{ad } g)^{d_2+1}v = 0$, for some d_1, d_2 such that $d_1 + d_2 = d$. The space of elements of finite order in V is a $(\mathbf{H}_c\text{-}\mathbf{H}_{c'})$ -sub-bimodule, to be denoted by V_{fin} . Clearly, if V is a finitely generated bimodule, then V_{fin} is the maximal Harish-Chandra subbimodule in V . For example, $\mathcal{D}(X_c)_{\text{fin}} = \mathcal{D}(X_c)$.

Further, given a *left* $\mathbf{H}_c$ -module M , and a *left* $\mathbf{H}_{c'}$ -module N , the vector space $\text{Hom}_{\mathbb{C}}(M, N)$ has a canonical $(\mathbf{H}_c\text{-}\mathbf{H}_{c'})$ -bimodule structure, and we put $\text{Hom}_{\text{fin}}(M, N) := (\text{Hom}_{\mathbb{C}}(M, N))_{\text{fin}}$. Similar definitions apply to $(\mathbf{eH}_c\mathbf{e}\text{-}\mathbf{eH}_{c'}\mathbf{e})$ -bimodules.

Since we will be dealing with standard modules over Cherednik algebras with varying parameter ‘ c ’, it will be convenient to incorporate this parameter in the notation and to write $M(\tau, c)$ for the standard module over $\mathbf{H}_c$, and $M_{\mathbf{e}}(\tau, c) := \mathbf{e} \cdot M(\tau, c)$ for the corresponding standard module over $\mathbf{eH}_c\mathbf{e}$. Recall that triv denotes the trivial 1-dimensional representation of the group W .

For any $\tau \in \text{Irrep}(W)$ and $c, c' \in \mathbb{C}[R]^W$, we define an object ${}_c\mathbf{P}_{c'}(\tau) \in \mathcal{HC}_{c,c'}(\mathbf{eHe})$ by

$${}_c\mathbf{P}_{c'}(\tau) := \text{Hom}_{\text{fin}}(M_{\mathbf{e}}(\tau^*, c'), M_{\mathbf{e}}(\text{triv}, c)). \quad (8.9)$$

Proposition 8.10. (i) *For any $c \in \mathbb{C}[R]_{\text{reg}}^W$, the $\mathbf{eH}_c\mathbf{e}$ -action on $M_{\mathbf{e}}(\text{triv}, c)$ induces a natural $\mathbf{eH}_c\mathbf{e}$ -bimodule isomorphism: $\mathbf{eH}_c\mathbf{e} \xrightarrow{\sim} {}_c\mathbf{P}_c(\text{triv})$.*

(ii) *The space ${}_c\mathbf{P}_{c'}(\text{triv})$ is nonzero if and only if $c - c'$ is an integer valued function.*

(iii) *For any $c, c', c'' \in \mathbb{C}[R]_{\text{reg}}^W$ such that $c - c', c' - c'' \in \mathbb{Z}[R]^W$, there is a canonical bimodule isomorphism*

$${}_c\mathbf{P}_{c'}(\text{triv}) \bigotimes_{\mathbf{eH}_{c'}\mathbf{e}} {}_{c'}\mathbf{P}_{c''}(\text{triv}) \xrightarrow{\sim} {}_c\mathbf{P}_{c''}(\text{triv}).$$

Proof. Part (i) is formulated here for the reader’s convenience only; it is a special case of the more general Theorem 8.15 formulated and proved below.

To prove (ii) we use the *shift operator* $\mathbf{S}_{c',c'+m}$, see [O] and also [EG2]. Suppose that $c - c'$ is integer valued, and let us show that ${}_c\mathbf{P}_{c'}(\text{triv})$ is nonzero. First assume that $c = c' + m$, where $m \in \mathbb{Z}[R]_{+}^W$ is nonnegative. Then the shift operator $\mathbf{S}_{c',c'+m}$ is a (nonzero) element of ${}_c\mathbf{P}_{c'}(\text{triv})$. Indeed, it is of “degree 0” with respect to the algebra of Calogero Hamiltonians, and of finite degree with respect to the algebra of functions (as it is a differential operator). On the other hand, if $m \leq 0$, i.e. if $(-m) \in \mathbb{Z}[R]_{+}^W$, then there also exists a shift operator $\mathbf{S}_{c',c'+m}$, which however does not preserve the space of symmetric polynomials (it has poles). Nevertheless, the operator $\delta(x)^{2N}\mathbf{S}_{c',c'+m}$ is a nonzero element of ${}_c\mathbf{P}_{c'}(\text{triv})$ for large enough N . Since any integer valued function m is a sum of a nonnegative and a nonpositive function, we find that ${}_c\mathbf{P}_{c'}(\text{triv})$ is nonzero for any integer valued $c - c'$.

Conversely, if ${}_c\mathbf{P}_{c'}(\text{triv}) \neq 0$ then it must have elements of degree 0 under the adjoint action of Calogero hamiltonians. Such an element satisfies the axiomatics of a shift operator of Opdam. But Opdam showed that shift operators exist only if $c - c'$ is integer: indeed, if a shift operator exists then the monodromies of the Calogero system for parameters c and c' are the same, i.e. $e^{2\pi ic} = e^{2\pi ic'}$. Part (ii) is proved.

To prove (iii), denote the left hand side of the isomorphism by H . It is clear that the action of any element $u \in H$ gives a map $\hat{u} : M_{\mathbf{e}}(\text{triv}, c'') \rightarrow M_{\mathbf{e}}(\text{triv}, c)$ such that $\hat{u} \in \text{Hom}_{\text{fin}}(M_{\mathbf{e}}(\text{triv}, c''), M_{\mathbf{e}}(\text{triv}, c))$. This way we obtain a natural bimodule homomorphism $j : H \rightarrow {}_c\mathbf{P}_{c''}(\text{triv})$. Localizing at the subset $\mathfrak{h}^{\text{reg}}/W \subset \mathfrak{h}/W$ we get a homomorphism $j|_{\mathfrak{h}^{\text{reg}}/W} : H|_{\mathfrak{h}^{\text{reg}}/W} \rightarrow {}_c\mathbf{P}_{c''}(\text{triv})|_{\mathfrak{h}^{\text{reg}}/W}$. The latter morphism is independent of the choice of the parameter ‘ c ’, and is easily seen to be bijective

for $c = 0$. Hence $j|_{\mathfrak{h}_{\text{reg}}/W}$ is a bijection. Next, we use an analogue of Corollary 3.6 (for bimodules over Cherednik algebras with two possibly different parameters ‘ c ’). The proof of this result is entirely analogous to the proof of Corollary 3.6. It follows from the result that the kernel of j has no torsion. We conclude that j is injective. Similarly, the cokernel of j has no torsion by the same analogue of Corollary 3.6. Thus, j is surjective, hence bijective. $\square$

Remark. The map $\mathbf{e}H_c\mathbf{e} \rightarrow \text{Hom}_{\text{fin}}(M_{\mathbf{e}}(\text{triv}, c), M_{\mathbf{e}}(\text{triv}, c))$ of Proposition 8.10 is also an *algebra* isomorphism with respect to the composition structure on the Hom -space on the right.

We now give an alternative, stronger and more effective version of Morita equivalence of Harish-Chandra categories that arises from Morita equivalence of algebras $\mathbf{e}H_c\mathbf{e}$ and $\mathbf{e}H_{c-m}\mathbf{e}$ provided by Theorem 8.1.

Proposition 8.11. *Let $c' \in \mathbb{C}[R]_{\text{reg}}^W$ and $c = c' + m$, where $m \in \mathbb{Z}[R]_+^W$.*

- (i) *The equivalence of categories $\mathcal{HC}_c(\mathbf{e}H\mathbf{e})$ and $\mathcal{HC}_{c'}(\mathbf{e}H\mathbf{e})$ arising from Morita equivalence of algebras $\mathbf{e}H_c\mathbf{e}$ and $\mathbf{e}H_{c'}\mathbf{e}$ is given by the functor: $V \mapsto {}_cP_c(\text{triv}) \otimes_{\mathbf{e}H_c\mathbf{e}} V \otimes_{\mathbf{e}H_{c'}\mathbf{e}} {}_{c'}P_{c'}(\text{triv})$.*
- (ii) *Moreover, this functor is an equivalence of monoidal categories.*

Proof. Clearly, it suffices to consider the case $m = \mathbf{1}_\varepsilon$. In that case the Morita equivalence of algebras $\mathbf{e}H_c\mathbf{e}$ and $\mathbf{e}H_{c'}\mathbf{e}$ comes from the identification $\mathbf{e}_\varepsilon H_c \mathbf{e}_\varepsilon = \mathbf{e}H_{c-\mathbf{1}_\varepsilon}\mathbf{e}$ of Proposition 4.11. We will use an isomorphism of bimodules: $\mathbf{e}H_c\mathbf{e} \simeq \text{Hom}_{\text{fin}}(\mathbf{e}_\varepsilon \cdot M(\text{triv}, c), \mathbf{e} \cdot M(\text{triv}, c))$, which is proved similarly to the case where $\mathbf{e}_\varepsilon$ is replaced by $\mathbf{e}$, see Proposition 8.10. Further, it is easy to see that under the identification $\mathbf{e}_\varepsilon H_c \mathbf{e}_\varepsilon = \mathbf{e}H_{c-\mathbf{1}_\varepsilon}\mathbf{e}$ of Proposition 4.11, the module $\mathbf{e}_\varepsilon \cdot M(\text{triv}, c)$ goes to $\mathbf{e} \cdot M(\text{triv}, c - \mathbf{1}_\varepsilon)$. So, for $m = \mathbf{1}_\varepsilon$, the equivalence of Harish-Chandra categories arising from Theorem 8.1 is provided by the bimodule $\text{Hom}_{\text{fin}}(\mathbf{e} \cdot M(\text{triv}, c - m), \mathbf{e} \cdot M(\text{triv}, c))$ which is exactly the bimodule ${}_cP_{c'}(\text{triv})$. Part (i) follows.

To prove (ii) one has to verify the isomorphism: ${}_cP_{c'}(\text{triv}) \otimes_{\mathbf{e}H_{c'}\mathbf{e}} {}_{c'}P_c(\text{triv}) = {}_cP_c(\text{triv})$, as well as an associativity constraint:

$${}_cP_{c'}(\text{triv}) \bigotimes_{\mathbf{e}H_{c'}\mathbf{e}} {}_{c'}P_{c''}(\text{triv}) \xrightarrow{\sim} {}_cP_{c''}(\text{triv}).$$

The first isomorphism follows from the equality $\mathbf{e}H_c\mathbf{e}_\varepsilon H_c\mathbf{e} = \mathbf{e}H_c\mathbf{e}$, by simplicity of the algebra $\mathbf{e}H_c\mathbf{e}$, and the second is part (iii) of Proposition 8.10. $\square$

Let $\varepsilon : W \rightarrow \{\pm 1\}$ be a character. Then the algebra H_c is isomorphic to $H_{\varepsilon c}$. Thus, we have established the ‘*if*’ part of the following

Conjecture 8.12. Let $c \in \mathbb{C}[R]_{\text{reg}}^W$. Then the algebras $\mathbf{e}H_c\mathbf{e}$ and $\mathbf{e}H_{c'}\mathbf{e}$, resp. H_c and $H_{c'}$, are Morita equivalent if and only if there exists a character $\varepsilon : W \rightarrow \{\pm 1\}$ such that $c - \varepsilon c' \in \mathbb{Z}[R]^W$.

For $\mathfrak{h} = \mathbb{C}$ and $W = \mathbb{Z}/2\mathbb{Z}$, the conjecture is a known result due to Hodges [Ho]. Also, for $W = S_n$ and transcendental c , this conjecture has been proved in [BEG2].

Recall that if we write $\mathbb{K} = \mathbb{C}(\mathfrak{h})$ for the field of rational functions on $\mathfrak{h}$, then we have $\mathcal{D}(X_c) = \{u \in \mathcal{D}(\mathbb{K}) \mid u(Q_c) \subset Q_c\}$, see (7.2). Motivated by this, for any $c, c' \in \mathbb{Z}[R]_+^W$, we put $\mathcal{D}(Q_{c'}, Q_c) := \{u \in \mathcal{D}(\mathbb{K}) \mid u(Q_{c'}) \subset Q_c\}$. The space $\mathcal{D}(Q_{c'}, Q_c)$ has a natural right $\mathcal{D}(X_{c'})$ -module structure, and left $\mathcal{D}(X_c)$ -module structure. Generalizing the proof of simplicity of the algebra $\mathcal{D}(X_c)$ one obtains

Proposition 8.13. *The $\mathcal{D}(X_c)$ - $\mathcal{D}(X_{c'})$ -bimodule $\mathcal{D}(Q_{c'}, Q_c)$ is simple, $\forall c, c' \in \mathbb{Z}[R]_+^W$.* $\square$

We can now consider the vector space $\mathcal{K}_{c,c'}$ of "kernels" with parameters c, c' . Specifically, we put

$$\mathcal{K}_{c,c'} = \left\{ P \in \mathbb{C}[\mathfrak{h} \times \mathfrak{h}^*] \mid \begin{array}{l} (x, k) \mapsto P(x, k) \cdot e^{(x,k)} \text{ is } c\text{-quasi-invariant with} \\ \text{respect to } x, \text{ and } c'\text{-quasi-invariant with respect to } k \end{array} \right\}.$$

Similarly to the case $c = c'$, one proves

Proposition 8.14. *For any $c, c' \in \mathbb{Z}[R]_+^W$,*

(i) *The space $\mathcal{K}_{c,c'}$ is a rank one projective $Q_c \otimes Q_{c'}$ -module.*

(ii) *There is a natural $Q_c \otimes Q_{c'}$ -module isomorphism: $\mathcal{D}(Q_{c'}, Q_c) \xrightarrow{\sim} \mathcal{K}_{c,c'}$.* $\square$

Further, the group W acts naturally on $\mathcal{D}(Q_{c'}, Q_c)$ commuting with $(\mathbf{e}H_c \mathbf{e} - \mathbf{e}H_{c'} \mathbf{e})$ -bimodule structure. This gives a W -module and $(\mathbf{e}H_c \mathbf{e} - \mathbf{e}H_{c'} \mathbf{e})$ -bimodule decomposition

$$\mathcal{D}(Q_{c'}, Q_c) = \bigoplus_{\tau \in \text{Irrep}(W)} \tau^* \otimes D_{c,c'}(\tau) \quad \text{where} \quad D_{c,c'}(\tau) := (\mathcal{D}(Q_{c'}, Q_c) \otimes \tau)^W.$$

Theorem 8.15. (i) *For any $\sigma, \tau \in \text{Irrep}(W)$, there is a natural $(\mathbf{e}H_c \mathbf{e} - \mathbf{e}H_{c'} \mathbf{e})$ -bimodule isomorphism*

$$\text{Hom}_{\text{fin}}(M_{\mathbf{e}}(\sigma'_{c'}, c'), M_{\mathbf{e}}(\tau'_c, c)) = \bigoplus_{\xi \in \text{Irrep}(W)} \text{Hom}_W(\xi \otimes \sigma, \tau) \otimes D_{c,c'}(\xi).$$

(ii) *The $(\mathbf{e}H_c \mathbf{e} - \mathbf{e}H_{c'} \mathbf{e})$ -bimodules $D_{c,c'}(\tau)$, $\tau \in \text{Irrep}(W)$, are irreducible, and pairwise non-isomorphic.*

Proof. To prove (i), denote the right hand side of the isomorphism by V and the left hand side by V' . Then $V = (\mathcal{D}(Q_{c'}, Q_c) \otimes \text{Hom}_{\mathbb{C}}(\sigma, \tau))^W$. On the other hand, by Proposition 6.6, $M_{\mathbf{e}}(\tau'_c) = (\tau \otimes Q_c)^W$. For any $v \in \text{Hom}_W(\xi \otimes \sigma, \tau) \otimes (\mathcal{D}(Q_{c'}, Q_c) \otimes \xi)^W \subset V$, the natural action-map $v : M_{\mathbf{e}}(\sigma'_{c'}, c') \rightarrow M_{\mathbf{e}}(\tau'_c, c)$ gives rise to a bimodule morphism $\theta : V \rightarrow V'$. The morphism θ is injective since, for any $v \in V$, $v \neq 0$, the induced action-map $v : M_{\mathbf{e}}(\sigma'_{c'}, c')|_{\mathfrak{h}^{\text{reg}}/W} \rightarrow M_{\mathbf{e}}(\tau'_c, c)|_{\mathfrak{h}^{\text{reg}}/W}$ is given by a differential operator, which is clearly nonzero. Thus, V may (and will) be viewed as a subbimodule of V' .

Now, consider the localized bimodules $V|_{\mathfrak{h}^{\text{reg}}/W}$ and $V'|_{\mathfrak{h}^{\text{reg}}/W}$ over the localized algebra $\mathbf{e}H_c \mathbf{e}|_{\mathfrak{h}^{\text{reg}}/W}$. We have: $\mathbf{e}H_c \mathbf{e}|_{\mathfrak{h}^{\text{reg}}/W} \simeq \mathcal{D}(\mathfrak{h})^W|_{\mathfrak{h}^{\text{reg}}/W}$, and $V|_{\mathfrak{h}^{\text{reg}}/W} \simeq (\mathcal{D}(\mathfrak{h}) \otimes \text{Hom}_{\mathbb{C}}(\sigma, \tau))^W|_{\mathfrak{h}^{\text{reg}}/W}$. It is clear (from nilpotency of the adjoint $\mathbb{C}[\mathfrak{h}]^W$ -action on V') that any element of $V'|_{\mathfrak{h}^{\text{reg}}/W}$ is represented by a differential operator, so $V'|_{\mathfrak{h}^{\text{reg}}/W} = V|_{\mathfrak{h}^{\text{reg}}/W}$. Thus, $(V'/V)|_{\mathfrak{h}^{\text{reg}}/W} = (V'|_{\mathfrak{h}^{\text{reg}}/W})/(V|_{\mathfrak{h}^{\text{reg}}/W}) = 0$, and $V'/V = 0$, by an analogue of Corollary 4.5. Part (i) is proved. Proof of part (ii) is entirely analogous to the proof of Theorem 7.17. $\square$

Remark. Under the equivalence of Theorem 8.5, the algebra $\mathcal{D}(X_c)$ corresponds to an algebra A in the tensor category $\text{Rep}(W)$ isomorphic to the regular representation of W . It is easy to show that in fact $A = \text{Fun}(W)$.

Recall the locally finite $SL(2)$ -action on $\mathcal{D}(X_c)$. When we pass to $\mathbf{gr}(\mathcal{D}(X_c))$, this action extends to a $GL(2)$ -action, since the order operator is added to the Lie algebra.

Question. What is the $GL(2)$ -character of each isotypic component of $\mathbf{gr}(\mathcal{D}(X_c))$?

This question is equivalent to the problem of computing the two-variable Hilbert series of isotypic components $\mathbf{gr}(\mathcal{D}(X_c))$, with respect to gradings both in $\mathfrak{h}$ and in $\mathfrak{h}^*$. Also, by the above results, it is equivalent to the question of computing the "characters" of all Harish-Chandra modules. It would be interesting to use the bijection $\mathcal{D}(X_c) \xrightarrow{\sim} \mathcal{K}_c$ in order to find the character of $\mathbf{gr}(\mathcal{D}(X_c))$.

Since $\mathbf{e}H_c \mathbf{e}$ is Morita equivalent to H_c , there is an $\mathbf{e}H_c \mathbf{e}$ -counterpart of Theorem 8.15. Specifically, one defines H_c -bimodules $V(\tau) := H_c \mathbf{e} \otimes_{\mathbf{e}H_c \mathbf{e}} D(\tau) \otimes_{\mathbf{e}H_c \mathbf{e}} \mathbf{e}H_c$ and proves

Theorem 8.16. *The bimodules $V(\tau)$ are irreducible and pairwise non-isomorphic. Moreover, for any $\sigma, \tau \in \text{Irrrep}(W)$, one has a H_c -bimodule isomorphism*

$$\text{Hom}_{\text{fin}}(M(\sigma'_c), M(\tau'_c)) = \bigoplus_{\xi \in \text{Irrrep}(W)} \text{Hom}_W(\xi \otimes \sigma, \tau) \otimes V(\xi). \quad \square$$

9 Applications of the shift operator

In this section we prove some further results about the structure of the variety X_c and the algebra $\mathcal{D}(X_c)$ using the notion of a *shift operator* introduced by Opdam. Recall that, given an *integral* c , a shift operator S_c is defined as a differential operator in $\mathcal{D}(\mathfrak{h})$ that satisfies the relation $L_c \circ S_c = S_c \circ L_0$ and has the principal symbol $\delta_c(x) \delta_c(k)$. We remark that the operator S_c determines (and is uniquely determined by) the Baker-Akhiezer function $\psi_c(x, k)$, introduced in §7, via the equation $\psi_c(x, k) = S_c[e^{(x, k)}]$. Below, however, we prefer to work with the shift operator rather than with $\psi_c(x, k)$ directly.

It follows from the results of [O] and [He] (see also [CV]) that such an operator exists and is unique for any $c \in \mathbb{Z}[R]_+^W$. In fact, S_c can be given explicitly by the following formula (cf. [Be], (2.11)):

$$S_c = \frac{1}{2^d d!} \text{ad}(L_c, L_0)^d [\delta_c(x)^2], \quad (9.1)$$

where $d := \sum_{\alpha \in R_+} c_\alpha$ and $\text{ad}(L_c, L_0)$ denotes the adjoint action on $\mathcal{D}(\mathfrak{h}^{\text{reg}})$ defined by multiplying the elements of $\mathcal{D}(\mathfrak{h}^{\text{reg}})$ by L_c on the left and by L_0 on the right.

Our applications of the shift operator S_c will be based on the following observation due to Opdam [O] (see also [He], (1.21)).

Lemma 9.2. $S_c[1] \in \mathbb{C} \setminus \{0\}$ for any $c \in \mathbb{Z}[R]_+^W$.

First, we use S_c to give another proof of bijectivity of normalization $\pi_c : \tilde{X}_c \rightarrow X_c$ (cf. Lemma 7.3(ii)):

For an arbitrary smooth affine variety X , and $\mathfrak{m}_1, \mathfrak{m}_2$, two distinct maximal ideals of $A = \mathbb{C}[X]$, we have:

$$\text{Hom}_A(A/\mathfrak{m}_1^k, A/\mathfrak{m}_2) = 0 \quad \text{for all } k = 1, 2, \dots \quad (9.3)$$

Now, fix $c \in \mathbb{Z}[R]_+^W$. By Lemma 6.3, we have $\tilde{X}_c \cong \mathfrak{h}$. Hence we may (and will) identify $\mathbb{C}[\tilde{X}_c] = \mathbb{C}[\mathfrak{h}]$, and write for short $A := \mathbb{C}[\mathfrak{h}]$ and $B := \mathbb{C}[X_c]$. Thus, $\tilde{X}_c = \text{Spec } A$, $X_c = \text{Spec } B$, and the normalization map $\pi_c : \text{Spec } A \rightarrow \text{Spec } B$ is given by $\mathfrak{m} \mapsto \mathfrak{m} \cap B$. Assume that π_c is *not* injective. Then there are (at least) two distinct maximal ideals $\mathfrak{m}_1$ and $\mathfrak{m}_2$ in A , such that $\mathfrak{m}_1 \cap B = \mathfrak{m}_2 \cap B$. Let S_c be the shift operator introduced above. It is easy to see from formula (9.1) (and Lemma 6.4) that $\text{order}(S_c) = d$ and $S_c \in \mathcal{D}(A, B)$. Hence $S_c[\mathfrak{m}_1^k] \subset \mathfrak{m}_1$ for all $k > d$, and therefore $S_c[\mathfrak{m}_1^k] \subseteq \mathfrak{m}_1 \cap B = \mathfrak{m}_2 \cap B \subseteq \mathfrak{m}_2$. It follows that, for each $k > d$, S_c induces a differential operator $\bar{S}_c \in \mathcal{D}(A/\mathfrak{m}_1^k, A/\mathfrak{m}_2)$. But, if $\mathfrak{m}_1 \neq \mathfrak{m}_2$, (9.3) means that $\mathcal{D}_0(A/\mathfrak{m}_1^k, A/\mathfrak{m}_2) = 0$ and then $\mathcal{D}(A/\mathfrak{m}_1^k, A/\mathfrak{m}_2) = 0$ by the inductive definition of differential operators (see (7.1)). Thus $\bar{S}_c = 0$, which implies that $S_c[A] \subseteq \mathfrak{m}_2$. In particular, since $1 \in A$, we get $S_c[1] \in \mathfrak{m}_2$, with obvious contradiction to Lemma 9.2. The result follows. $\square$

Now, let X be an affine algebraic variety with smooth normalization $\pi : \tilde{X} \rightarrow X$. Since $\mathbb{C}[X] \subseteq \mathbb{C}[\tilde{X}]$, we may set

$$\mathcal{D}(\tilde{X}, X) := \{ u \in \mathcal{D}(\tilde{X}) \mid u(\mathbb{C}[\tilde{X}]) \subseteq \mathbb{C}[X] \}. \quad (9.4)$$

Then $\mathcal{D}(\tilde{X}, X)$ is a right ideal in $\mathcal{D}(\tilde{X})$ and (with identifications (7.2)) a left ideal of $\mathcal{D}(X)$. According to [SS], Lemma 2.7, $\mathcal{D}(\tilde{X}, X)$ is isomorphic to the space $\mathcal{D}(\mathbb{C}[\tilde{X}], \mathbb{C}[X])$ of all differential operators from $\mathbb{C}[\tilde{X}]$ to $\mathbb{C}[X]$ over $\mathbb{C}[X]$, so the notation (9.4) is consistent with (7.1). We write $\text{End}_{\mathcal{D}(\tilde{X})} \mathcal{D}(\tilde{X}, X)$ for the endomorphism ring of $\mathcal{D}(\tilde{X}, X)$ as a right $\mathcal{D}(\tilde{X})$ -module and identify it with a subalgebra in $\mathcal{Q}$, the quotient skew-field of $\mathcal{D}(\tilde{X})$ (cf. §10 below).

Theorem 9.5. $\mathcal{D}(X_c) = \text{End}_{\mathcal{D}(\tilde{X}_c)} \mathcal{D}(\tilde{X}_c, X_c)$.

Proof. Clearly, $\mathcal{D}(X_c) \subseteq \text{End}_{\mathcal{D}(\tilde{X}_c)} (\mathcal{D}(\tilde{X}_c, X_c))$. Hence, we only need to show the opposite inclusion. To this end, observe first that $\mathcal{D}(\tilde{X}_c, X_c)(\mathbb{C}[\tilde{X}_c]) = \mathbb{C}[X_c]$. Indeed, by definition we have $\mathcal{D}(\tilde{X}_c, X_c)(\mathbb{C}[\tilde{X}_c]) \subseteq \mathbb{C}[X_c]$. On the other hand, by Lemma 9.2 and the fact that $S_c \in \mathcal{D}(\tilde{X}_c, X_c)$, we get

$$\mathbb{C}[X_c] = \mathbb{C}[X_c] \circ S_c[1] \subseteq \mathcal{D}(\tilde{X}_c, X_c)[1] \subseteq \mathcal{D}(\tilde{X}_c, X_c)(\mathbb{C}[\tilde{X}_c]).$$

Now, write $\mathbb{K} = \mathbb{C}(X_c) \cong \mathbb{C}(\mathfrak{h})$ for the field of rational functions on X_c . Then, $\text{End}_{\mathcal{D}(\tilde{X}_c)} (\mathcal{D}(\tilde{X}_c, X_c)) \subset \mathcal{D}(\mathbb{K})$ by (10.1). Further, for any $u \in \text{End}_{\mathcal{D}(\tilde{X}_c)} \mathcal{D}(\tilde{X}_c, X_c)$, we have

$$u(\mathbb{C}[X_c]) = u \circ \mathcal{D}(\tilde{X}_c, X_c)(\mathbb{C}[\tilde{X}_c]) \subseteq \mathcal{D}(\tilde{X}_c, X_c)(\mathbb{C}[\tilde{X}_c]) = \mathbb{C}[X_c].$$

Hence $u \in \mathcal{D}(X_c)$. □

Next, we recall a well-known ring-theoretic result usually referred to as the "Dual Basis Lemma". Let A be an algebra, $\mathcal{P}$ a right A -module, and $B = \text{End}_A \mathcal{P}$, the endomorphism ring of $\mathcal{P}$, and $\mathcal{P}^* = \text{Hom}_A(\mathcal{P}, A)$ its dual module. Write $\mathcal{P} \cdot \mathcal{P}^*$ for the subspace of $\text{End}_A(\mathcal{P})$ spanned by the endomorphisms $p \cdot f : x \mapsto p \cdot f(x)$, for all $p \in \mathcal{P}$ and $f \in \mathcal{P}^* = \text{Hom}_A(\mathcal{P}, A)$. Obviously $\mathcal{P} \cdot \mathcal{P}^*$ is a two-sided ideal of $B = \text{End}_A(\mathcal{P})$. Similarly, $\mathcal{P}^* \cdot \mathcal{P}$ stands for a two-sided ideal of A generated by the elements $\{f(q) \mid q \in \mathcal{P} \text{ and } f \in \mathcal{P}^*\}$.

Lemma 9.6 ([MR], 3.5.2). *The A -module $\mathcal{P}$ is finitely generated and projective if and only if $\mathcal{P} \cdot \mathcal{P}^* = B$.* □

Now we are in a position to prove one of the main results in the present paper.

Theorem 9.7. *The ring $\mathcal{D}(X_c)$ is Morita equivalent to $\mathcal{D}(\mathfrak{h})$.*

Proof. In the setup of Lemma 9.6, write $A := \mathcal{D}(\tilde{X}_c)$, $B := \mathcal{D}(X_c)$, and $\mathcal{P} := \mathcal{D}(\tilde{X}_c, X_c)$. Then, by Theorem 9.5, we have $B = \text{End}_A \mathcal{P}$. Now, $\mathcal{P} \cdot \mathcal{P}^*$ is a nonzero two-sided ideal of B , and therefore, is equal to B by Theorem 7.20. By Lemma 9.6, we then conclude that $\mathcal{P}$ is finitely generated projective. On the other hand, $\mathcal{P}^* \cdot \mathcal{P}$ is a nonzero two-sided ideal of $A \cong \mathcal{D}(\mathfrak{h})$, and hence, by simplicity of the latter, is equal to A . This means that $\mathcal{P}$ is a generator in the category of (right) A -modules. Summing up, we have $B = \text{End}_A \mathcal{P}$, where $\mathcal{P}$ is a finitely generated projective generator in $\text{mod-}A$. By the Morita Theorem, A and B are then equivalent rings. □

The Morita equivalence of Theorem 9.7 yields

Corollary 9.8. *The homological dimension of the algebra $\mathcal{D}(X_c)$ equals $\dim \mathfrak{h}$.* □

We end up this section with a conjecture giving an alternative description of the category of modules over $\mathcal{D}(X_c)$ hinted by [DE].

Let $i : X_c \hookrightarrow Y$ be a closed imbedding of X_c into a *smooth* affine algebraic variety Y . Write $\mathcal{D}(X, Y) := \mathcal{D}(\mathbb{C}[Y], \mathbb{C}[X_c])$ for the space of differential operators from $\mathbb{C}[Y]$ to $\mathbb{C}[X_c]$ over $\mathbb{C}[Y]$. Let $\text{mod-}\mathcal{D}(X_c)$ denote the abelian category of right $\mathcal{D}(X_c)$ -modules, and $\text{mod}_{X_c}\text{-}\mathcal{D}(Y)$ denote the abelian category of right $\mathcal{D}(Y)$ -modules with support in $i(X_c) \subset Y$.

Conjecture 9.9. The direct image functor i_+ is an equivalence of categories, with quasi-inverse $i^+ : \text{mod}_{X_c}\text{-}\mathcal{D}(Y) \longrightarrow \text{mod-}\mathcal{D}(X_c)$, $N \longmapsto \text{Hom}_{\mathcal{D}(Y)}(\mathcal{D}(X, Y), N)$.

Conjecture 9.9 can be viewed as a generalization of a well-known result of Kashiwara to the *singular* variety X_c . In case when $\dim X_c = 1$ (or more generally, for any singular affine curve with injective normalization) this conjecture has been proved in [DE].

Conjecture 9.9 has been proved in full generality in [BN].

10 Appendix: A filtration on differential operators

The goal of this section is to put the filtration (7.12) into a more general context of the ring of differential operators on a (singular) algebraic variety X whose normalisation is a vector space, cf. also [P] in the special case $\dim X = 1$.

Let X be any irreducible (not necessarily smooth) algebraic variety, $\pi : \tilde{X} \rightarrow X$ its normalisation, and $\mathbb{K} = \mathbb{C}(X) = \mathbb{C}(\tilde{X})$, the field of rational functions. The ring $\mathcal{D}(\mathbb{K})$ is a Noetherian domain (see, e.g., [MR], 15.5.5). By Goldie's Theorem, it has the quotient skew-field which we denote by $\mathcal{Q}$. Let $\mathcal{P} := \mathcal{D}(\tilde{X}, X)$, regarded as a right ideal of $\mathcal{D}(\tilde{X})$, see (9.4). We put $\mathcal{E} := \text{End}_{\mathcal{D}(\tilde{X})} \mathcal{P}$, the endomorphism ring of $\mathcal{P}$. Then we may (and will) identify $\mathcal{E} \cong \{q \in \mathcal{Q} : q\mathcal{P} \subseteq \mathcal{P}\}$ (see [MR], 3.1.15). Since $\mathcal{P}$ is also a left ideal of $\mathcal{D}(X)$, we have $\mathcal{D}(X) \subseteq \mathcal{E}$. On the other hand, $\mathbb{C}[\tilde{X}]$ being finite over $\mathbb{C}[X]$ implies that $f \cdot \mathbb{C}[\tilde{X}] \subseteq \mathbb{C}[X]$ for some nonzero $f \in \mathbb{C}[\tilde{X}]$, hence $\mathcal{P} \cap \mathbb{C}[\tilde{X}] \neq \{0\}$ and therefore $\mathcal{P} \in \mathcal{D}(\mathbb{K}) \Rightarrow \mathcal{E} \subset \mathcal{D}(\mathbb{K})$. We summarize these inclusions in the following

Lemma 10.1 (see [SS]). $\mathcal{P} \subset \mathcal{D}(X) \subseteq \mathcal{E} \subset \mathcal{D}(\mathbb{K})$, $\mathcal{P} \subset \mathcal{D}(\tilde{X}) \subset \mathcal{D}(\mathbb{K})$. $\square$

By definition, the algebra $\mathcal{D}(\mathbb{K})$ carries a natural (differential) filtration. Using (10.1), we may (and will) equip the spaces $\mathcal{P}, \mathcal{E}, \mathcal{D}(X), \mathcal{D}(\tilde{X})$ with the induced filtrations⁶. We write $\text{gr } \mathcal{P}, \text{gr } \mathcal{E}, \text{gr } \mathcal{D}(X), \text{gr } \mathcal{D}(\tilde{X})$ for the associated graded objects. The next result is probably known to the experts, but we cannot find a precise reference in the literature.

Proposition 10.2. *Let X be an irreducible affine algebraic variety over $\mathbb{C}$. Assume that the normalization $\tilde{X}$ is smooth. Then*

$$\text{gr } \mathcal{P} \subset \text{gr } \mathcal{D}(X) \subseteq \text{gr } \mathcal{E} \subseteq \text{gr } \mathcal{D}(\tilde{X}) \subset \text{gr } \mathcal{D}(\mathbb{K}) . \quad (10.3)$$

Proof. Given the inclusions (10.1), we need only to prove $\text{gr } \mathcal{E} \subseteq \text{gr } \mathcal{D}(\tilde{X})$. First, recall that $\tilde{X}$ being smooth implies that $\text{gr } \mathcal{D}(\tilde{X}) \cong \mathbb{C}[T^*\tilde{X}]$ is a regular Noetherian commutative domain. Hence $\text{gr } \mathcal{P} \subseteq \text{gr } \mathcal{D}(\tilde{X})$ is a finitely generated ideal of $\text{gr } \mathcal{D}(\tilde{X})$. Since $\mathbb{C}[\tilde{X}]$ is finite over $\mathbb{C}[X]$, $\mathcal{P}$ is nonzero: in fact, $\mathcal{P} \supset \text{Ann}_{\mathbb{C}[X]}(\mathbb{C}[\tilde{X}]/\mathbb{C}[X]) \neq \{0\}$. Next, with our description of the endomorphism ring $\mathcal{E}$, we have $\mathcal{E} \cdot \mathcal{P} \subseteq \mathcal{P}$ in $\mathcal{D}(\mathbb{K})$, and therefore $(\text{gr } \mathcal{E}) \cdot (\text{gr } \mathcal{P}) \subseteq \text{gr } \mathcal{P}$ in $\text{gr } \mathcal{D}(\mathbb{K})$. Hence, by [AM], Prop. 2.4, the elements of $\text{gr } \mathcal{E}$ are integral over $\text{gr } \mathcal{D}(\tilde{X})$ in $\text{gr } \mathcal{D}(\mathbb{K})$. But, being regular, the ring $\text{gr } \mathcal{D}(\tilde{X})$ is integrally closed, hence $\text{gr } \mathcal{E} \subseteq \text{gr } \mathcal{D}(\tilde{X})$. $\square$

Remark. If $\dim X = 1$ then $\tilde{X}$ is always smooth. In this case our Proposition 10.2 implies Proposition 3.11 of [SS] which establishes the inclusion $\text{gr } \mathcal{D}(X) \subseteq \text{gr } \mathcal{D}(\tilde{X})$ for any (irreducible affine) curve.

⁶The induced filtrations on $\mathcal{D}(X)$ and $\mathcal{D}(\tilde{X})$ coincide with the intrinsic ones under identifications (7.2).

Now assume that X is an irreducible variety with normalization $\tilde{X}$ being isomorphic to V , a vector space. We identify $\mathbb{C}(X)$ with $\mathbb{K} = \mathbb{C}(V)$. On $\mathbb{K}$, introduce an increasing filtration $\{F_j^\flat \mathbb{K}\}_{j \in \mathbb{Z}}$ as follows. Let $\mathbb{P}(V \oplus \mathbb{C}) \supset V$ be the projective completion of V , and $D_\infty = \mathbb{P}(V \oplus \mathbb{C}) \setminus V$, the divisor at infinity. Given $f \in \mathbb{K} = \mathbb{C}(V)$, we can think of f as a rational function on $\mathbb{P}(V \oplus \mathbb{C})$, and we put $f \in F_j^\flat \mathbb{K}$ if f has a pole of order $\leq j$ at D_∞ . The restriction of this filtration on $\mathbb{K}$ to $\mathbb{C}[V]$ coincides with the standard filtration by degree of polynomial. We extend the filtration $F_j^\flat \mathbb{K}$ to the algebra $\mathcal{D}(\mathbb{K})$ of differential operators with rational coefficients by assigning filtration degree zero to all constant coefficient differential operators. We write $\{\mathcal{D}_n^\flat(\mathbb{K})\}_{n \in \mathbb{Z}}$ for the induced filtration on $\mathcal{D}(\mathbb{K})$ and call it the $\flat$ -filtration. Now, if E is a linear subspace of $\mathcal{D}(\mathbb{K})$, we set $E_n^\flat := E \cap \mathcal{D}_n^\flat(\mathbb{K})$ and write $\mathrm{gr}^\flat E := \bigoplus_{n \in \mathbb{Z}} E_n^\flat / E_{n-1}^\flat$ for the associated graded space. Clearly, by symmetry with the standard filtration, we have $\mathrm{gr}^\flat \mathcal{D}(\tilde{X}) \cong \mathbb{C}[V \times V^*]$, and therefore $\mathrm{gr}^\flat \mathcal{D}(\mathbb{K})$ is isomorphic to a (properly graded) localization of $\mathbb{C}[V \times V^*]$ at $\mathbb{C}[V] \setminus \{0\}$. In particular, both $\mathrm{gr}^\flat \mathcal{D}(\tilde{X})$ and $\mathrm{gr}^\flat \mathcal{D}(\mathbb{K})$ are commutative domains, the former being regular. Reviewing the proof of Proposition 10.2 then shows that its argument works for the $\flat$ -filtration as well. Thus, with the above notation and conventions, we have the following

Proposition 10.4. *If X is an irreducible affine variety with normalization $\tilde{X}$ being isomorphic to V , then $\mathrm{gr}^\flat \mathcal{P} \subset \mathrm{gr}^\flat \mathcal{D}(X) \subseteq \mathrm{gr}^\flat \mathcal{E} \subseteq \mathrm{gr}^\flat \mathcal{D}(\tilde{X}) \subset \mathrm{gr}^\flat \mathcal{D}(\mathbb{K})$.*

By definition, the differential operators of order zero form a maximal commutative, (locally) ad-nilpotent subalgebra of $\mathcal{D}(X)$, which is naturally isomorphic to $\mathbb{C}[X]$. On the other hand, from Proposition 10.4 we obtain the following generalization of the result (proved in §6) on the nilpotency of the adjoint action on $\mathcal{D}(X_c)$ of the subalgebra $Q_c^\flat = \mathcal{D}_0^\flat(X_c)$.

Proposition 10.5. *If $\tilde{X} \cong V$, then $\mathcal{D}_0^\flat(X)$ is a commutative (locally) ad-nilpotent subalgebra of $\mathcal{D}(X)$.*

Proof. Since $\mathrm{gr}^\flat \mathcal{D}(X) \subset \mathbb{C}[V \times V^*]$, the elements of $\mathcal{D}(X)$ have non-negative $\flat$ -degrees. Thus $\mathcal{D}_{-1}^\flat(X) = \{0\}$. In view of commutativity of $\mathrm{gr}^\flat \mathcal{D}(\mathbb{K})$, we have $[\mathcal{D}_0^\flat(X), \mathcal{D}_n^\flat(X)] \subset \mathcal{D}_{n-1}^\flat(X)$ for all $n \geq 0$. Hence $\mathcal{D}_0^\flat(X)$ is ad-nilpotent. Moreover, if $n = 0$, we have $[\mathcal{D}_0^\flat(X), \mathcal{D}_0^\flat(X)] \subseteq \mathcal{D}_{-1}^\flat(X) = \{0\}$, therefore $\mathcal{D}_0^\flat(X)$ is commutative. $\square$

References

- [AM] M. F. Atiyah and I. G. Macdonald, *Introduction to Commutative Algebra*, Addison-Wesley Publishing Co., Reading, 1969.
- [BW] Yu. Berest and G. Wilson, *Classification of rings of differential operators on affine curves*, Internat. Math. Res. Notices **2** (1999), 105–109.
- [Be] Yu. Berest, *Huygens' principle and the bispectral problem* in *The Bispectral Problem*, CRM Proceedings and Lecture Notes, **14**, Amer. Math. Soc. 1998, pp. 11–30.
- [BEG1] Y. Berest, P. Etingof, and V. Ginzburg, Finite-dimensional representations of rational Cherednik algebras, Int. Math. Res. Not. (2003), no. 19, pp. 1053–1088.
- [BEG2] Y. Berest, P. Etingof, and V. Ginzburg, Morita equivalence of Cherednik algebras, J. Reine Angew. Math., 568 (2004), 8198.
- [BGG] J. Bernstein, I. Gelfand and S. Gelfand, *A certain category of $\mathfrak{g}$ -modules*. (Russian) Funct. Anal. Appl. 10:2 (1976), 1–8.
- [BGG1] J. Bernstein, I. Gelfand and S. Gelfand, *Differential operators on the cubic cone*, Russian Math. Surveys **27** (1972), 466–488.
- [B] J. E. Björk, *Rings of Differential Operators*, North-Holland Mathematical Library **21**, North-Holland Publishing Co., Amsterdam, 1979.
- [BMR] M. Broué, G. Malle and R. Rouquier, *Complex reflection groups, braid groups, Hecke algebras*, J. Reine Angew. Math. **500** (1998), 127–190.

- [BN] D. Ben-Zvi, T. Nevins, *Cusps and $\mathcal{D}$ -modules*. J. Amer. Math. Soc. **17** (2004), 155–179.
- [C] O. A. Chalykh, *Darboux transformations for multidimensional Schrödinger operators*, Russian Math. Surveys **53**(2) (1998), 377–379.
- [CV] O. A. Chalykh and A. P. Veselov, *Commutative rings of partial differential operators and Lie algebras*, Comm. Math. Phys. **126**(3) (1990), 597–611.
- [CV2] O. A. Chalykh and A. P. Veselov, *Integrability in the theory of Schrödinger operator and harmonic analysis*, Comm. Math. Phys. **152**(1) (1993), 29–40.
- [CS] M. Chamarie and J. T. Stafford, *When rings of differential operators are maximal orders*, Math. Proc. Camb. Phil. Soc. **102** (1987), 399–410.
- [Ch] I. Cherednik, *Lectures on affine KZ equations, quantum many-body problems, Hecke algebras, and Macdonald theory*, Lecture Notes by Date et al., Kyoto, 1995.
- [Ch2] I. Cherednik, *Double affine Hecke algebras, Knizhnik-Zamolodchikov equations, and Macdonald operators*, IMRN (Duke math.J.) **9** (1992), p.171–180.
- [CG] N. Chriss, V. Ginzburg, *Representation theory and complex geometry*. Birkhäuser Boston, 1997.
- [Dz] C. Dezelee, *Représentations de dimension finie de l’algèbre de Cherednik rationnelle*. [arXiv:math.RT/0111210](#).
- [DE] M. Van Doorn and A. Van den Essen, *$\mathcal{D}$ -Modules with support on a curve*, Publ. RIMS **23** (1987), 937–953.
- [Di] J. Dixmier, *Enveloping algebras*. Graduate Studies in Mathem. **11**, A.M.S., Providence, RI, 1996.
- [D] C. F. Dunkl, *Differential-difference operators and monodromy representations of Hecke algebras*, Pacific J. Math. **159**(2) (1993), 271–298.
- [DO] C. F. Dunkl and E. Opdam, *Dunkl operators for complex reflection groups*, [arXiv:math.RT/0108185](#).
- [DJO] C. F. Dunkl, M. F. E. de Jeu and E. Opdam, *Singular polynomials for finite reflection groups*, Trans. Amer. Math. Soc. **346** (1994), 237–256.
- [EG] P. Etingof and V. Ginzburg, *Symplectic reflection algebras, Calogero-Moser space, and deformed Harish-Chandra homomorphism*. Invent. Math. **147** (2002), 243–348, [[arXiv:math.AG/0011114](#)].
- [EG2] P. Etingof and V. Ginzburg, *On m -quasi-invariants of Coxeter groups*, Preprint [arXiv:math.QA/0106175](#), to appear in Moscow Mathem. Journ.
- [FV] M. Feigin and A. Veselov, *Quasi-invariants of Coxeter groups and m -harmonic polynomials*, Int. Math. Res. Not. **10** (2002), 521–545. [[arXiv:math-ph/0105014](#)].
- [FeV] G. Felder and A. Veselov, *Action of Coxeter groups on m -harmonic polynomials and KZ equations*, Preprint 2001, [arXiv:QA/0108012](#).
- [Gi] V. Ginzburg, *On primitive ideals*, [arXiv:math.RT/0202079](#).
- [Go] I. Gordon, *Baby Verma modules for rational Cherednik algebras*. [arXiv:math.RT/0202301](#).
- [G] A. Grothendieck, *Eléments de Géométrie Algébrique IV*, Publ. Math. **32**, IHES, Paris, 1967.
- [Gu] N. Guay, *Projectives in the category $\mathcal{O}$ for the Cherednik algebra*. Preprint, University of Chicago 2001.
- [GGOR] V. Ginzburg, N. Guay, E. Opdam, R. Rouquier, *On the category $\mathcal{O}$ for rational Cherednik algebras*, Invent. Math. **154** (2003), no. 3, 617–651.
- [He] G. Heckman, *A Remark on the Dunkl differential-difference operators*. Harmonic analysis on reductive groups, 181–191, Progr. Math., **101**, Birkhäuser Boston, Boston, MA, 1991.
- [Ho] T. Hodges, *Morita equivalence of primitive factors of $U(\mathfrak{sl}_2)$* , Contemp. Math. **139** (1992), 175–179.
- [HS] R. Hart and S. P. Smith, *Differential operators on some singular surfaces*, Bull. London Math. Soc. **19** (1987), 145–148.
- [K] V. Kac, *Constructing groups associated to infinite dimensional Lie algebras*, in: Infinite dimensional groups and appl., MSRI Publ. **4**, Springer-Verlag, 1985, 167–216.
- [Ku] A. Kuznetsov, *Quiver varieties and Hilbert schemes*. [arXiv:math.AG/0111092](#)
- [LS] T. Levasseur and J. Stafford, *Invariant differential operators and a homomorphism of Harish-Chandra*, J. Amer. Math. Soc. **8** (1995), 365–372.
- [LS2] T. Levasseur and J.T. Stafford, *Semi-simplicity of invariant holonomic systems on a reductive Lie algebra*. Amer. J. Math. **119** (1997), 1095–1117.
- [Ma] I.G.Macdonald, *Symmetric functions and Hall polynomials*, second edition, Oxford University Press (1995).
- [MR] J. C. McConnell and J. C. Robson, *Noncommutative Noetherian Rings*, J.Wiley & Sons, NY 1987.

- [Mo] S. Montgomery, *Fixed Rings of Finite Automorphism Groups of Associative Rings*, LN in Math., **818** Springer-Verlag, Berlin/New-York, 1980.
- [Mu] I. M. Musson, *Some rings of differential operators which are Morita equivalent to the Weyl algebra*, Proc. Amer. Math. Soc. **98** (1986), 29–30.
- [MvdB] I. M. Musson, M. Van den Bergh, *Invariants under tori of rings of differential operators and related topics*. Mem. Amer. Math. Soc. **136** (1998), no. 650.
- [O] E. Opdam, *Dunkl operators, Bessel functions and the discriminant of a finite Coxeter group*, Compositio Math. **85**(3) (1993), 333–373.
- [O1] E. M. Opdam, *A remark on the irreducible characters and fake degrees of finite real reflection groups*, Invent. Math. **120** (3) (1995), 447–454.
- [OR] E. Opdam and R. Rouquier, private communication, July 2001.
- [P] P. Perkins, *Commutative subalgebras of the ring of differential operators on a curve*, Pacific Journal of Math. **139** (1989), 279–302.
- [Sm] S. P. Smith, *An example of a ring Morita equivalent to the Weyl algebra*, J. Algebra **73** (1981), 552–555.
- [SS] S. P. Smith and J. T. Stafford, *Differential operators on an affine curve*, Proc. London Math. Soc. (3) **56** (1988), 229–259.
- [St] J. T. Stafford, *Homological properties of the enveloping algebra $U(\mathfrak{sl}_2)$* , Math. Proc. Cambr. Phil. Soc. **91** (1982), 29–37.
- [S] R. Stanley, *Invariants of finite groups and their applications to combinatorics*, Bull. Amer. Math. Soc. (N.S.) **1**(3) (1979), 475–511.
- [S2] R. Stanley, *Enumerative combinatorics. Vol. 2*. Cambridge Studies in Advanced Mathematics, **62** Cambridge University Press, Cambridge, 1999.
- [VdB] M. Van den Bergh, *Differential operators on semi-invariants for tori and weighted projective spaces in Topics in Invariant Theory*, Lecture Notes in Math. **1478**, Springer, Berlin, 1991, pp. 255–272.
- [VSC] A. P. Veselov, K. L. Styrkas and O. A. Chalykh, *Algebraic integrability for the Schrödinger equation, and groups generated by reflections*, Theoret. and Math. Phys. **94**(2) (1993), 182–197.
- [Wa] N. Wallach, *Invariant differential operators on a reductive Lie algebra and Weyl group representations*, J. Amer. Math. Soc. **6** (1993), 779–816.
- [Wi] G. Wilson, *Collisions of Calogero-Moser particles and an adelic Grassmannian*, Invent. Mathem. **133** (1998), 1–41.

Y.B.: Department of Mathematics, Cornell University, Ithaca, NY 14853-4201, USA;
berest@math.cornell.edu

P.E.: Department of Mathematics, Rm 2-165, MIT, 77 Mass. Ave, Cambridge, MA 02139;
etingof@math.mit.edu

V.G.: Department of Mathematics, University of Chicago, Chicago, IL 60637, USA;
ginzburg@math.uchicago.edu